










among infants with alive at discharge




among infants with alive at discharge









among infants with followup at 6 years of age




among infants with followup at 6 years of age




among infants with home medical care




among infants with home medical care




among infants with home medical care




among infants with home medical care




among infants with home medical care



among infants with home medical care




among infants with home medical care




among infants with home medical care







among infants with home medical care




among infants with home medical care




among infants with followup at 6 years of age




among infants with followup at 6 years of age




among infants with visual impairment




among infants with visual impairment




among infants with followup at 6 years of age




among infants with followup at 6 years of age




among infants with followup at 6 years of age




among infants with followup at 6 years of age




among infants with hearing impairment




among infants with hearing impairment




among infants with hearing impairment



among infants with hearing impairment




among infants with followup at 6 years of age




among infants with followup at 6 years of age







among infants with motor impairment





among infants with motor impairment




among infants with cerebral palsy




among infants with cerebral palsy







among infants with cerebral palsy




among infants with cerebral palsy




among infants with cerebral palsy




among infants with cerebral palsy




among infants with followup at 6 years of age




among infants with followup at 6 years of age




among infants with followup at 6 years of age



among infants with followup at 6 years of age




among infants with followup at 6 years of age




among infants with followup at 6 years of age








among infants with IQ or DQ by other methods




among infants with IQ or DQ by other methods











among infants with followup at 6 years of age




among infants with followup at 6 years of age










among infants with followup at 6 years of age



among infants with followup at 6 years of age