







among infants with alive at discharge





among infants with alive at discharge




among infants with alive at discharge




among infants with alive at discharge







among infants with followup at 6 years of age




among infants with followup at 6 years of age




among infants with home medical care




among infants with home medical care




among infants with home medical care




among infants with home medical care




among infants with home medical care




among infants with home medical care




among infants with home medical care





among infants with home medical care




among infants with home medical care




among infants with home medical care







among infants with visual impairment




among infants with visual impairment




among infants with followup at 6 years of age




among infants with followup at 6 years of age




among infants with followup at 6 years of age





among infants with followup at 6 years of age




among infants with hearing impairment




among infants with hearing impairment








among infants with followup at 6 years of age




among infants with followup at 6 years of age







among infants with cerebral palsy




among infants with cerebral palsy




among infants with cerebral palsy




among infants with cerebral palsy




among infants with cerebral palsy




among infants with cerebral palsy




among infants with cerebral palsy




among infants with cerebral palsy




among infants with followup at 6 years of age





among infants with followup at 6 years of age




among infants with followup at 6 years of age




among infants with followup at 6 years of age








among infants with IQ or DQ by other methods




among infants with IQ or DQ by other methods







among infants with followup at 6 years of age




among infants with followup at 6 years of age




among infants with followup at 6 years of age





among infants with followup at 6 years of age




among infants with followup at 6 years of age




among infants with followup at 6 years of age





among infants with followup at 6 years of age



among infants with followup at 6 years of age