



among infants with number of fetus 2>



Plurality





among infants with number of fetus 2>





Diabetes





Pregnancy induced hypertension



Clinical CAM




Histologic CAM




among infants with positive histologic CAM



Grade of histologic CAM





among infants with positive histologic CAM





PROM





Maternal steroid





NRFS





Presentation





Mode of delivery





Cord blood transfusion



Gender




Neonatal transport




among infants with inborn



Maternal transport





among infants with inborn





Oxygen use at birth



Intubation at birth




Live birth




among infants with live birth and remained






among infants with live birth and remained




among infants with live birth and remained

Air leak syndrome




among infants with live birth and remained




among infants with live birth and remained


Pulmonary hemorrhage




among infants with live birth and remained




among infants with live birth, remained and mechanical ventilation






among infants with live birth, remained and mechanical ventilation




among infants with live birth, remained and alive at 28 days of age

CLD at 28 d




among infants with live birth, remained and alive at 28 days of age




among infants with CLD


Glucocorticoid for CLD




among infants with CLD




among infants with live birth, remained, alive at 36 wk(corrected age), and CLD






among infants with live birth, remained, alive at 36 wk(corrected age), and CLD




among infants with live birth and remained

PDA with symptom




among infants with live birth and remained




among infants with live birth and remained

Indomethacin for PDA




among infants with live birth and remained




among infants with symptomatic PDA

Surgical ligation for PDA




among infants with symptomatic PDA




among infants with live birth, remained and alive at 7 d

Late onset adrenal insufficiency




among infants with live birth, remained and alive at 7 d




among infants with live birth and remained

Seizure




among infants with live birth and remained




among infants with live birth and remained

Intraventricular hemorrhage




among infants with live birth and remained




among infants with live birth, remained and IVH


Post IVH hydrocephalus




among infants with live birth, remained and IVH




among infants with live birth and remained






among infants with live birth and remained




among infants with live birth and remained

HIE




among infants with live birth and remained




among infants with live birth and remained

Intrauterine infection




among infants with live birth and remained




among infants with live birth and remained

Sepsis




among infants with live birth and remained




among infants with live birth and remained

Use of antibiotics




among infants with live birth and remained




among infants with live birth and remained

Intravenous hyperalimentation




among infants with live birth and remained




among infants with live birth and remained

NEC




among infants with live birth and remained




among infants with live birth and remained

Idiopathic intestinal perforation




among infants with live birth and remained




among infants with live birth and remained

Hearing loss screening




among infants with live birth and remained




among infants with live birth and remained



Treatment for ROP



among infants with live birth and remained




Congenital anomaly




among infants with live birth, remained and congenital anomaly

Operation for congenital anomaly




among infants with live birth, remained and congenital anomaly




among infants with live birth and remained

Blood transfusion




among infants with live birth and remained




among infants with live birth and remained

Erythropoietin




among infants with live birth and remained




among infants with live birth and remained

Dead at discharge




among infants with live birth and remained




among infants with live birth

Dead at discharge




among infants with live birth




among infants with live birth, remained and dead at discharge

Autopsy




among infants with live birth, remained and dead at discharge




among infants with live birth, remained and alive at discharge

Discharge home




among infants with live birth, remained and alive at discharge




among infants with live birth, remained and alive at discharge

HOT




among infants with live birth, remained and alive at discharge



among infants with live birth and alive at discharge

Tracheostomy



among infants with live birth and alive at discharge