

























No.	Resources of participating hospitals	All hospitals	n
<b>R</b>			
<b>2410</b>	Followup at 6 years of age (among infants with alive at discharge)	 <p>1:Yes 3% 2:No 97%</p>	65174
<b>2412</b>	Dead after discharge (among infants with alive at discharge)	 <p>1:Yes 1% 2:No 46% 3:not available 53%</p>	1117
<b>2416</b>	Reason for dropout (among infants with alive at discharge)	 <p>1:Followed at different hospital 44% 2:Admitted in rehabilitaion center 0% 3:No contact 41% 4:Others 14%</p>	1033
<b>2420</b>	Age at followup (mean) (among infants with followup at 6 years of age)	5.8	2248
	SD	0.4	
	95% confidence interval	5.8-5.8	
<b>2422</b>	Age corrected at followup (mean) (among infants with followup at 6 years of age)	5.3	2148
	SD	0.5	
	95% confidence interval	5.3-5.3	
<b>2430</b>	Body weight (mean) (among infants with followup at 6 years of age)	18.0	2180
	SD	3.1	
	95% confidence interval	17.9-18.1	

No.	Resources of participating hospitals	All hospitals	n
2440	Height (mean) (among infants with followup at 6 years of age)	109.5	2165
	SD	5.4	
	95% confidence interval	109.3-109.7	
2450	Head circumference (mean) (among infants with followup at 6 years of age)	50.2	1686
	SD	1.9	
	95% confidence interval	50.1-50.3	
2452	Chest circumference (mean) (among infants with followup at 6 years of age)	53.5	1179
	SD	3.8	
	95% confidence interval	53.2-53.7	
2454	Abdominal circumference (mean) (among infants with followup at 6 years of age)	50.6	204
	SD	4.7	
	95% confidence interval	50.0-51.3	
2455	Systolic blood pressure (mean) (among infants with followup at 6 years of age)	102.5	445
	SD	11.1	
	95% confidence interval	101.5-103.6	
2456	Diastolic blood pressure (mean) (among infants with followup at 6 years of age)	61.6	444
	SD	10.1	
	95% confidence interval	60.6-62.5	

No.	Resources of participating hospitals	All hospitals	n
2460	Home oxygen use (among infants with home medical care)	 <p>1:Yes 53% 2:No 47%</p>	58
2550	Home medical care (among infants with followup at 6 years of age)	 <p>1:Yes 4% 2:No 96%</p>	2124
2551	Mechanical ventilation (among infants with home medical care)	 <p>1:Yes 11% 2:No 89%</p>	38
2552	Tracheostomy (among infants with home medical care)	 <p>1:Yes 15% 2:No 85%</p>	40
2553	Tube feeding (among infants with home medical care)	 <p>1:Yes 49% 2:No 51%</p>	47
2554	VP shunt (among infants with home medical care)	 <p>1:Yes 36% 2:No 64%</p>	39

No.	Resources of participating hospitals	All hospitals	n
2560	Duration of home oxygen (mean) (among infants with followup at 6 years of age)	20.8	67
	SD	14.6	
	95% confidence interval	17.3-24.3	
2570	Visual impairment (among infants with followup at 6 years of age)	 <p>1:Yes 14% 2:No 86%</p>	2115
2572	Severity of visual impairment (among infants with visual impairment)	 <p>1:Less than light perception 2% 2:Finger movement, nystagmus 1% 3:Amblyopia 58% 4:Strabismus 39% 5:Others 0% 6:Unknown 0%</p>	170
2574	Eye glasses (among infants with followup at 6 years of age)	 <p>1:Yes 13% 2:No 87%</p>	2015
2580	Hearing impairment (among infants with followup at 6 years of age)	 <p>1:Yes 2% 2:No 98%</p>	2129
2582	Hearing aid (among infants with hearing impairment)	 <p>1:Yes 62% 2:No 38%</p>	40







No.	Resources of participating hospitals	All hospitals	n
2584	Hearing level with aid (among infants with hearing impairment)	 <p>1:No 24% 2:Moderate 52% 3:Severe 24%</p>	25
2600	Motor impairment (among infants with followup at 6 years of age)	 <p>1:Yes 9% 2:No 91%</p>	2141
2610	Level of motor impairment (among infants with motor impairment)	 <p>1:Mild 20% 2:Cerebral palsy 80%</p>	172
2620	GMFCS grade (among infants with cerebral palsy)	 <p>1: I 22% 2: II 20% 3: III 16% 4: IV 16% 5: V 27%</p>	90
2630	Type of cerebral palsy (among infants with cerebral palsy)	 <p>1:Spastic 82% 2:Athetoid 7% 3:Mixed 2% 4:Flaccid 6% 5:Others 2%</p>	125
2640	Topographical distribution (among infants with cerebral palsy)	 <p>1:Quadriplegia 37% 2:Diplegia 29% 3:Paraplegia 18% 4:Hemiplegia 13% 5:Monoplegia 2%</p>	120

No.	Resources of participating hospitals	All hospitals	n
2650	Cause of cerebral palsy (among infants with cerebral palsy)	 <p>A bar chart with a vertical axis from 0 to 2500 in increments of 500. The horizontal axis has three categories: EYL, IVH, and Others. The bar for EYL reaches 1950, the bar for IVH reaches 294, and the bar for Others is very low, near 0.</p>	120
2660	DQ or IQ measurement (among infants with followup at 6 years of age)	 <p>A pie chart with two segments. Segment 1 is the larger portion, labeled '1' and '87%'. Segment 2 is the smaller portion, labeled '2' and '13%'.</p>	2244
2670	Reason not to measure DQ or IQ (among infants with followup at 6 years of age)	 <p>A pie chart with six segments labeled 1 through 6. Segment 1 is the largest, followed by segment 2. Segments 3, 4, and 5 are very small, while segment 6 is slightly larger than them.</p>	250
2680	Method for IQ or DQ measurement (among infants with followup at 6 years of age)	 <p>A pie chart with seven segments labeled 1 through 7. Segment 1 is the dominant portion. Segments 2, 3, 4, 5, 6, and 7 are all very small slices.</p>	1945
2700	WISCIV Full IQ (mean) (among infants with IQ by WISCIV)	92.3	1492
	SD	14.3	
	95% confidence interval	91.6-93.0	
2702	WISCIV VCI (Verbal) (mean) (among infants with IQ by WISCIV)	95.5	1495
	SD	13.3	
	95% confidence interval	94.8-96.1	

No.	Resources of participating hospitals	All hospitals	n
2704	WISCIV PRI (Perceptual) (mean) (among infants with IQ by WISCIV)	94.7	1497
	SD	14.9	
	95% confidence interval	93.9-95.4	
2706	WISCIV WMI (Working) (mean) (among infants with IQ by WISCIV)	90.4	1496
	SD	16.3	
	95% confidence interval	89.6-91.2	
2708	WISCIV PSI (Processing) (mean) (among infants with IQ by WISCIV)	92.0	1490
	SD	14.9	
	95% confidence interval	91.2-92.7	
2710	WPPSI Full IQ (mean) (among infants with IQ by WPPSI)	74.6	29
	SD	21.2	
	95% confidence interval	66.8-82.3	
2712	WPPSI VCI (Verbal) (mean) (among infants with IQ by WPPSI)	69.6	26
	SD	18.3	
	95% confidence interval	62.6-76.6	
2714	WPPSI PRI (Perceptual) (mean) (among infants with IQ by WPPSI)	80.7	24
	SD	23.0	
	95% confidence interval	71.5-89.9	

No.	Resources of participating hospitals	All hospitals	n
2716	WPPSI PSI (Processing) (mean) (among infants with IQ by WIPPSI)	73.9	16
	SD	23.2	
	95% confidence interval	62.6-85.3	
2718	WPPSI GLC (Global language) (mean) (among infants with IQ by WIPPSI)	68.8	9
	SD	17.4	
	95% confidence interval	57.4-80.2	
2820	New Kyoto scale (mean) (among infants with DQ by K scale)	66.3	165
	SD	25.3	
	95% confidence interval	62.4-70.1	
2830	Tanaka-Binet scale (mean) (among infants with IQ by T-B)	79.0	91
	SD	20.3	
	95% confidence interval	74.9-83.2	
2840	K-ABCII scale (mean) (among infants with IQ by K-ABC)	0.0	0
	SD	0.0	
	95% confidence interval	0.0-0.0	
2850	Other method (among infants with IQ or DQ by other methods)	<p>1:Normal 76% 2:Boderline 18% 3:Delay 6%</p>	121



No.	Resources of participating hospitals	All hospitals	n
2860	Evaluated by physician (among infants with followup at 6 years of age)	 <p>1:Normal 64% 2:Borderline 20% 3:Delay 15%</p>	1280
2870	Asthma (among infants with followup at 6 years of age)	 <p>1:Yes 7% 2:No 93%</p>	2089
2880	Epilepsy (among infants with followup at 6 years of age)	 <p>1:Yes 2% 2:No 98%</p>	2150
2890	Behavioral disorder (among infants with followup at 6 years of age)	 <p>1:Yes 15% 2:No 85%</p>	2108
2892	Type of behavioral disorder (among infants with behavior disorder)	 <p>ADHD: 591 ADHD...: 1535 ASD ASD... Others</p>	304
2900	Rehabilitation (among infants with followup at 6 years of age)	 <p>1:Yes 28% 2:No 72%</p>	2126