

























No.	Resources of participating hospitals	All hospitals	n
R			
2410	Followup at 6 years of age (among infants with alive at discharge)	 <p>1:Yes 4% 2:No 96%</p>	72915
2412	Dead after discharge (among infants with alive at discharge)	 <p>1:Yes 1% 2:No 49% 3:not available 50%</p>	1371
2416	Reason for dropout (among infants with alive at discharge)	 <p>1:Followed at different hospital 43% 2:Admitted in rehabilitation center 0% 3:No contact 42% 4:Others 15%</p>	1277
2420	Age at followup (mean) (among infants with followup at 6 years of age)	0.0	0
	SD	0.0	
	95% confidence interval	0.0-0.0	
2422	Age corrected at followup (mean) (among infants with followup at 6 years of age)	0.0	0
	SD	0.0	
	95% confidence interval	0.0-0.0	
2430	Body weight (mean) (among infants with followup at 6 years of age)	0.0	0
	SD	0.0	
	95% confidence interval	0.0-0.0	

No.	Resources of participating hospitals	All hospitals	n
2440	Height (mean) (among infants with followup at 6 years of age)	0.0	0
	SD	0.0	
	95% confidence interval	0.0-0.0	
2450	Head circumference (mean) (among infants with followup at 6 years of age)	0.0	0
	SD	0.0	
	95% confidence interval	0.0-0.0	
2452	Chest circumference (mean) (among infants with followup at 6 years of age)	0.0	0
	SD	0.0	
	95% confidence interval	0.0-0.0	
2454	Abdominal circumference (mean) (among infants with followup at 6 years of age)	0.0	0
	SD	0.0	
	95% confidence interval	0.0-0.0	
2455	Systolic blood pressure (mean) (among infants with followup at 6 years of age)	0.0	0
	SD	0.0	
	95% confidence interval	0.0-0.0	
2456	Diastolic blood pressure (mean) (among infants with followup at 6 years of age)	0.0	0
	SD	0.0	
	95% confidence interval	0.0-0.0	

No.	Resources of participating hospitals	All hospitals	n
2460	Home oxygen use (among infants with home medical care)	 <p>1: Yes 55% 2: No 45%</p>	78
2550	Home medical care (among infants with followup at 6 years of age)	 <p>1: Yes 4% 2: No 96%</p>	2753
2551	Mechanical ventilation (among infants with home medical care)	 <p>1: Yes 11% 2: No 89%</p>	56
2552	Tracheostomy (among infants with home medical care)	 <p>1: Yes 16% 2: No 84%</p>	58
2553	Tube feeding (among infants with home medical care)	 <p>1: Yes 42% 2: No 58%</p>	66
2554	VP shunt (among infants with home medical care)	 <p>1: Yes 32% 2: No 68%</p>	59







No.	Resources of participating hospitals	All hospitals	n
2560	Duration of home oxygen (mean) (among infants with followup at 6 years of age)	0.0	0
	SD	0.0	
	95% confidence interval	0.0-0.0	
2570	Visual impairment (among infants with followup at 6 years of age)	 <p>1:Yes 14% 2:No 86%</p>	2739
2572	Severity of visual impairment (among infants with visual impairment)	 <p>1:Less than light perception 2% 2:Finger movement, nystagmus 1% 3:Amblyopia 57% 4:Strabismus 40% 5:Others 0% 6:Unknown 0%</p>	230
2574	Eye glasses (among infants with followup at 6 years of age)	 <p>1:Yes 14% 2:No 86%</p>	2635
2580	Hearing impairment (among infants with followup at 6 years of age)	 <p>1:Yes 2% 2:No 98%</p>	2751
2582	Hearing aid (among infants with hearing impairment)	 <p>1:Yes 54% 2:No 46%</p>	50

No.	Resources of participating hospitals	All hospitals	n
2584	Hearing level with aid (among infants with hearing impairment)	 <p>1:No 24% 2:Moderate 52% 3:Severe 24%</p>	29
2600	Motor impairment (among infants with followup at 6 years of age)	 <p>1:Yes 9% 2:No 91%</p>	2778
2610	Level of motor impairment (among infants with motor impairment)	 <p>1:Mild 20% 2:Cerebral palsy 80%</p>	233
2620	GMFCS grade (among infants with cerebral palsy)	 <p>1: I 24% 2: II 21% 3: III 14% 4: IV 16% 5: V 25%</p>	122
2630	Type of cerebral palsy (among infants with cerebral palsy)	 <p>1:Spastic 83% 2:Athetoid 5% 3:Mixed 2% 4:Flaccid 7% 5:Others 2%</p>	166
2640	Topographical distribution (among infants with cerebral palsy)	 <p>1:Quadriplegia 41% 2:Diplegia 26% 3:Paraplegia 19% 4:Hemiplegia 12% 5:Monoplegia 3%</p>	158

No.	Resources of participating hospitals	All hospitals	n
2650	Cause of cerebral palsy (among infants with cerebral palsy)	 <p>A bar chart with a vertical axis from 0 to 3000 in increments of 500. The horizontal axis has three categories: PVL, IVH, and Others. The PVL bar reaches 2524, the IVH bar reaches 396, and the Others bar is at 0.</p>	164
2660	DQ or IQ measurement (among infants with followup at 6 years of age)	 <p>A pie chart with two segments. Segment 1 is the larger portion, labeled '1' and '86%'. Segment 2 is the smaller portion, labeled '2' and '14%'.</p>	2920
2670	Reason not to measure DQ or IQ (among infants with followup at 6 years of age)	 <p>A pie chart with six segments labeled 1 through 6. Segment 1 is the largest, followed by segment 6.</p>	318
2680	Method for IQ or DQ measurement (among infants with followup at 6 years of age)	 <p>A pie chart with seven segments labeled 1 through 7. Segment 1 is the dominant portion.</p>	2518
2700	WISCIV Full IQ (mean) (among infants with IQ by WISCIV)	0.0	0
	SD	0.0	
	95% confidence interval	0.0-0.0	
2702	WISCIV VCI (Verbal) (mean) (among infants with IQ by WISCIV)	0.0	0
	SD	0.0	
	95% confidence interval	0.0-0.0	

No.	Resources of participating hospitals	All hospitals	n
2704	WISCIV PRI (Perceptual) (mean) (among infants with IQ by WISCIV)	0.0	0
	SD	0.0	
	95% confidence interval	0.0-0.0	
2706	WISCIV WMI (Working) (mean) (among infants with IQ by WISCIV)	0.0	0
	SD	0.0	
	95% confidence interval	0.0-0.0	
2708	WISCIV PSI (Processing) (mean) (among infants with IQ by WISCIV)	0.0	0
	SD	0.0	
	95% confidence interval	0.0-0.0	
2710	WPPSI Full IQ (mean) (among infants with IQ by WIPPSI)	0.0	0
	SD	0.0	
	95% confidence interval	0.0-0.0	
2712	WPPSI VCI (Verbal) (mean) (among infants with IQ by WIPPSI)	0.0	0
	SD	0.0	
	95% confidence interval	0.0-0.0	
2714	WPPSI PRI (Perceptual) (mean) (among infants with IQ by WIPPSI)	0.0	0
	SD	0.0	
	95% confidence interval	0.0-0.0	

No.	Resources of participating hospitals	All hospitals	n
2716	WPPSI PSI (Processing) (mean) (among infants with IQ by WIPPSI)	0.0	0
	SD	0.0	
	95% confidence interval	0.0-0.0	
2718	WPPSI GLC (Global language) (mean) (among infants with IQ by WIPPSI)	0.0	0
	SD	0.0	
	95% confidence interval	0.0-0.0	
2820	New Kyoto scale (mean) (among infants with DQ by K scale)	0.0	0
	SD	0.0	
	95% confidence interval	0.0-0.0	
2830	Tanaka-Binet scale (mean) (among infants with IQ by T-B)	0.0	0
	SD	0.0	
	95% confidence interval	0.0-0.0	
2840	K-ABCII scale (mean) (among infants with IQ by K-ABC)	0.0	0
	SD	0.0	
	95% confidence interval	0.0-0.0	
2850	Other method (among infants with IQ or DQ by other methods)	<p>1:Normal 76% 2:Boderline 19% 3:Delay 6%</p>	135

No.	Resources of participating hospitals	All hospitals	n
2860	Evaluated by physician (among infants with followup at 6 years of age)	 <p>1: Normal 63% 2: Borderline 22% 3: Delay 15%</p>	1574
2870	Asthma (among infants with followup at 6 years of age)	 <p>1: Yes 8% 2: No 92%</p>	2720
2880	Epilepsy (among infants with followup at 6 years of age)	 <p>1: Yes 3% 2: No 97%</p>	2785
2890	Behavioral disorder (among infants with followup at 6 years of age)	 <p>1: Yes 16% 2: No 84%</p>	2727
2892	Type of behavioral disorder (among infants with behavior disorder)	 <p>ADHD: 767 ADHD...: 1977 ASD ASD... Others</p>	420
2900	Rehabilitation (among infants with followup at 6 years of age)	 <p>1: Yes 28% 2: No 72%</p>	2744