

























No.	Resources of participating hospitals	All hospitals	n
R			
2410	Followup at 6 years of age (among infants with alive at discharge)	 <p>1:Yes 5% 2:No 95%</p>	79624
2412	Dead after discharge (among infants with alive at discharge)	 <p>1:Yes 2% 2:No 53% 3:not available 45%</p>	1679
2416	Reason for dropout (among infants with alive at discharge)	 <p>1:Followed at different hospital 46% 2:Admitted in rehabilitation center 0% 3:No contact 39% 4:Others 15%</p>	1556
2420	Age at followup (mean) (among infants with followup at 6 years of age)	5.8	3726
	SD	0.4	
	95% confidence interval	5.8-5.8	
2422	Age corrected at followup (mean) (among infants with followup at 6 years of age)	5.3	3530
	SD	0.5	
	95% confidence interval	5.3-5.3	
2430	Body weight (mean) (among infants with followup at 6 years of age)	18.0	3630
	SD	3.0	
	95% confidence interval	17.9-18.1	

No.	Resources of participating hospitals	All hospitals	n
2440	Height (mean) (among infants with followup at 6 years of age)	109.5	3607
	SD	5.3	
	95% confidence interval	109.4-109.7	
2450	Head circumference (mean) (among infants with followup at 6 years of age)	50.3	2821
	SD	1.9	
	95% confidence interval	50.2-50.3	
2452	Chest circumference (mean) (among infants with followup at 6 years of age)	53.6	2007
	SD	3.7	
	95% confidence interval	53.4-53.8	
2454	Abdominal circumference (mean) (among infants with followup at 6 years of age)	50.8	583
	SD	4.7	
	95% confidence interval	50.4-51.2	
2455	Systolic blood pressure (mean) (among infants with followup at 6 years of age)	102.6	927
	SD	11.2	
	95% confidence interval	101.9-103.3	
2456	Diastolic blood pressure (mean) (among infants with followup at 6 years of age)	62.1	924
	SD	10.6	
	95% confidence interval	61.4-62.8	

No.	Resources of participating hospitals	All hospitals	n
2460	Home oxygen use (among infants with home medical care)	 <p>1: Yes 54% 2: No 46%</p>	93
2550	Home medical care (among infants with followup at 6 years of age)	 <p>1: Yes 4% 2: No 96%</p>	3505
2551	Mechanical ventilation (among infants with home medical care)	 <p>1: Yes 9% 2: No 91%</p>	70
2552	Tracheostomy (among infants with home medical care)	 <p>1: Yes 14% 2: No 86%</p>	72
2553	Tube feeding (among infants with home medical care)	 <p>1: Yes 39% 2: No 61%</p>	79
2554	VP shunt (among infants with home medical care)	 <p>1: Yes 35% 2: No 65%</p>	75

No.	Resources of participating hospitals	All hospitals	n
2560	Duration of home oxygen (mean) (among infants with followup at 6 years of age)	19.3	137
	SD	13.8	
	95% confidence interval	17.0-21.6	
2570	Visual impairment (among infants with followup at 6 years of age)	 <p>1:Yes 14% 2:No 86%</p>	3492
2572	Severity of visual impairment (among infants with visual impairment)	 <p>1:Less than light perception 2% 2:Finger movement, nystagmus 1% 3:Amblyopia 57% 4:Strabismus 40% 5:Others 0% 6:Unknown 0%</p>	270
2574	Eye glasses (among infants with followup at 6 years of age)	 <p>1:Yes 13% 2:No 87%</p>	3396
2580	Hearing impairment (among infants with followup at 6 years of age)	 <p>1:Yes 2% 2:No 98%</p>	3513
2582	Hearing aid (among infants with hearing impairment)	 <p>1:Yes 54% 2:No 46%</p>	59

No.	Resources of participating hospitals	All hospitals	n
2584	Hearing level with aid (among infants with hearing impairment)	 <p>1:No 23% 2: Moderate 54% 3: Severe 23%</p>	35
2600	Motor impairment (among infants with followup at 6 years of age)	 <p>1: Yes 9% 2: No 91%</p>	3535
2610	Level of motor impairment (among infants with motor impairment)	 <p>1: Mild 23% 2: Cerebral palsy 77%</p>	284
2620	GMFCS grade (among infants with cerebral palsy)	 <p>1: I 23% 2: II 24% 3: III 13% 4: IV 16% 5: V 24%</p>	146
2630	Type of cerebral palsy (among infants with cerebral palsy)	 <p>1: Spastic 83% 2: Athetoid 5% 3: Mixed 3% 4: Flaccid 7% 5: Others 3%</p>	192
2640	Topographical distribution (among infants with cerebral palsy)	 <p>1: Quadriplegia 40% 2: Diplegia 28% 3: Paraplegia 19% 4: Hemiplegia 11% 5: Monoplegia 2%</p>	178

No.	Resources of participating hospitals	All hospitals	n
2650	Cause of cerebral palsy (among infants with cerebral palsy)	 <p>A bar chart with a vertical axis from 0 to 3500. The first bar, labeled 'PVL', reaches 3264. The second bar, labeled 'IVH', reaches 453. The third bar, labeled 'Others', is very short and near zero.</p>	192
2660	DQ or IQ measurement (among infants with followup at 6 years of age)	 <p>A pie chart with two segments. Segment 1 is the large majority, labeled '1' and '88%'. Segment 2 is a small slice, labeled '2' and '12%'.</p>	3717
2670	Reason not to measure DQ or IQ (among infants with followup at 6 years of age)	 <p>A pie chart with six segments labeled 1 through 6. Segment 1 is the largest, followed by segment 6.</p>	361
2680	Method for IQ or DQ measurement (among infants with followup at 6 years of age)	 <p>A pie chart with eight segments labeled 1 through 8. Segment 1 is the largest, followed by segment 8.</p>	3233
2700	WISCIV Full IQ (mean) (among infants with IQ by WISCIV)	92.1	2454
	SD	14.4	
	95% confidence interval	91.5-92.6	
2702	WISCIV VCI (Verbal) (mean) (among infants with IQ by WISCIV)	95.2	2460
	SD	13.7	
	95% confidence interval	94.6-95.7	

Analysis results on infants born in 2003-2022

6 years of age XIII-7/11



No.	Resources of participating hospitals	All hospitals	n
2704	WISCIV PRI (Perceptual) (mean) (among infants with IQ by WISCIV)	95.0	2465
	SD	14.8	
	95% confidence interval	94.4-95.6	
2706	WISCIV WMI (Working) (mean) (among infants with IQ by WISCIV)	89.7	2462
	SD	16.5	
	95% confidence interval	89.0-90.3	
2708	WISCIV PSI (Processing) (mean) (among infants with IQ by WISCIV)	91.9	2454
	SD	14.7	
	95% confidence interval	91.3-92.5	
2710	WPPSI Full IQ (mean) (among infants with IQ by WIPPSI)	74.1	53
	SD	20.2	
	95% confidence interval	68.6-79.5	
2712	WPPSI VCI (Verbal) (mean) (among infants with IQ by WIPPSI)	72.3	50
	SD	18.6	
	95% confidence interval	67.2-77.5	
2714	WPPSI PRI (Perceptual) (mean) (among infants with IQ by WIPPSI)	81.1	48
	SD	19.6	
	95% confidence interval	75.6-86.7	

No.	Resources of participating hospitals	All hospitals	n
2716	WPPSI PSI (Processing) (mean) (among infants with IQ by WIPPSI)	75.1	40
	SD	20.3	
	95% confidence interval	68.8-81.4	
2718	WPPSI GLC (Global language) (mean) (among infants with IQ by WIPPSI)	67.2	16
	SD	14.4	
	95% confidence interval	60.1-74.2	
2720	WISCV Full scale IQ (mean) (among infants with IQ by WISCV)	0.0	0
	SD	0.0	
	95% confidence interval	0.0-0.0	
2722	WISCV VCI (Verbal) (mean) (among infants with IQ by WISCV)	0.0	0
	SD	0.0	
	95% confidence interval	0.0-0.0	
2724	WISCV VSI (Visual) (mean) (among infants with IQ by WISCV)	0.0	0
	SD	0.0	
	95% confidence interval	0.0-0.0	
2726	WISCV FRI (Fluid) (mean) (among infants with IQ by WISCV)	0.0	0
	SD	0.0	
	95% confidence interval	0.0-0.0	

No.	Resources of participating hospitals	All hospitals	n
2728	WISCV WMI (Working) (mean) (among infants with IQ by WISCV)	0.0	0
	SD	0.0	
	95% confidence interval	0.0-0.0	
2730	WISCV PSI (Processing) (mean) (among infants with IQ by WISCV)	0.0	0
	SD	0.0	
	95% confidence interval	0.0-0.0	
2732	WISCV QRI (Quantitative) (mean) (among infants with IQ by WISCV)	0.0	0
	SD	0.0	
	95% confidence interval	0.0-0.0	
2734	WISCV AWMI (Auditory) (mean) (among infants with IQ by WISCV)	0.0	0
	SD	0.0	
	95% confidence interval	0.0-0.0	
2736	WISCV NVI (Nonverbal) (mean) (among infants with IQ by WISCV)	0.0	0
	SD	0.0	
	95% confidence interval	0.0-0.0	
2738	WISCV GAI (General) (mean) (among infants with IQ by WISCV)	0.0	0
	SD	0.0	
	95% confidence interval	0.0-0.0	






Analysis results on infants born in 2003-2022

6 years of age XIII-10/11

No.	Resources of participating hospitals	All hospitals	n
2740	WISCV CPI (Cognitive) (mean) (among infants with IQ by WISCV)	0.0	0
	SD	0.0	
	95% confidence interval	0.0-0.0	
2820	New Kyoto scale (mean) (among infants with DQ by K scale)	65.2	268
	SD	23.9	
	95% confidence interval	62.3-68.1	
2830	Tanaka-Binet scale (mean) (among infants with IQ by T-B)	74.6	157
	SD	21.7	
	95% confidence interval	71.2-77.9	
2840	K-ABCII scale (mean) (among infants with IQ by K-ABC)	0.0	0
	SD	0.0	
	95% confidence interval	0.0-0.0	
2850	Other method (among infants with IQ or DQ by other methods)	 <p>1:Normal 73% 2:Boderline 21% 3:Delay 6%</p>	154
2860	Evaluated by physician (among infants with followup at 6 years of age)	 <p>1:Normal 62% 2:Boderline 22% 3:Delay 16%</p>	1906

Analysis results on infants born in 2003-2022

6 years of age XIII-11/11

No.	Resources of participating hospitals	All hospitals	n
2870	Asthma (among infants with followup at 6 years of age)	 <p>1: Yes 9% 2: No 91%</p>	3484
2880	Epilepsy (among infants with followup at 6 years of age)	 <p>1: Yes 2% 2: No 98%</p>	3550
2890	Behavioral disorder (among infants with followup at 6 years of age)	 <p>1: Yes 16% 2: No 84%</p>	3474
2892	Type of behavioral disorder (among infants with behavior disorder)	 <p>ADHD: 996 ADHD...: 2510 ASD ASD... Others</p>	548
2900	Rehabilitation (among infants with followup at 6 years of age)	 <p>1: Yes 28% 2: No 72%</p>	3506