

























No.	Resources of participating hospitals	All hospitals	n
R			
2410	Followup at 6 years of age (among infants with alive at discharge)	 1:Yes 6% 2:No 94%	86520
2412	Dead after discharge (among infants with alive at discharge)	 1:Yes 2% 2:No 57% 3:not available 41%	2432
2416	Reason for dropout (among infants with alive at discharge)	 1:Followed at different hospital 43% 2:Admitted in rehabilitation center 0% 3:No contact 36% 4:Others 21%	2270
2420	Age at followup (mean) (among infants with followup at 6 years of age)	5.8	5413
	SD	0.4	
	95% confidence interval	5.8-5.8	
2422	Age corrected at followup (mean) (among infants with followup at 6 years of age)	5.3	5130
	SD	0.5	
	95% confidence interval	5.3-5.3	
2430	Body weight (mean) (among infants with followup at 6 years of age)	17.9	5189
	SD	3.0	
	95% confidence interval	17.8-18.0	

No.	Resources of participating hospitals	All hospitals	n
2440	Height (mean) (among infants with followup at 6 years of age)	109.3	5150
	SD	5.4	
	95% confidence interval	109.1-109.4	
2450	Head circumference (mean) (among infants with followup at 6 years of age)	50.2	3933
	SD	2.0	
	95% confidence interval	50.2-50.3	
2452	Chest circumference (mean) (among infants with followup at 6 years of age)	53.6	2672
	SD	3.7	
	95% confidence interval	53.4-53.7	
2454	Abdominal circumference (mean) (among infants with followup at 6 years of age)	50.6	1021
	SD	4.4	
	95% confidence interval	50.3-50.9	
2455	Systolic blood pressure (mean) (among infants with followup at 6 years of age)	102.1	1434
	SD	11.2	
	95% confidence interval	101.5-102.7	
2456	Diastolic blood pressure (mean) (among infants with followup at 6 years of age)	61.9	1428
	SD	10.7	
	95% confidence interval	61.3-62.5	

No.	Resources of participating hospitals	All hospitals	n
2460	Home oxygen use (among infants with home medical care)	 1: Yes 49% 2: No 51%	140
2550	Home medical care (among infants with followup at 6 years of age)	 1: Yes 4% 2: No 96%	4905
2551	Mechanical ventilation (among infants with home medical care)	 1: Yes 14% 2: No 86%	116
2552	Tracheostomy (among infants with home medical care)	 1: Yes 18% 2: No 82%	117
2553	Tube feeding (among infants with home medical care)	 1: Yes 44% 2: No 56%	126
2554	VP shunt (among infants with home medical care)	 1: Yes 32% 2: No 68%	120

No.	Resources of participating hospitals	All hospitals	n
2560	Duration of home oxygen (mean) (among infants with followup at 6 years of age)	19.7	195
	SD	14.3	
	95% confidence interval	17.7-21.7	
2570	Visual impairment (among infants with followup at 6 years of age)	 1:Yes 13% 2:No 87%	4897
2572	Severity of visual impairment (among infants with visual impairment)	 1:Less than light perception 2% 2:Finger movement, nystagmus 1% 3:Amblyopia 58% 4:Strabismus 39% 5:Others 0% 6:Unknown 0%	374
2574	Eye glasses (among infants with followup at 6 years of age)	 1:Yes 13% 2:No 87%	4753
2580	Hearing impairment (among infants with followup at 6 years of age)	 1:Yes 2% 2:No 98%	4926
2582	Hearing aid (among infants with hearing impairment)	 1:Yes 50% 2:No 50%	86



No.	Resources of participating hospitals	All hospitals	n
2584	Hearing level with aid (among infants with hearing impairment)	 1:No 20% 2:Moderate 58% 3:Severe 22%	50
2600	Motor impairment (among infants with followup at 6 years of age)	 1:Yes 9% 2:No 91%	4944
2610	Level of motor impairment (among infants with motor impairment)	 1:Mild 25% 2:Cerebral palsy 75%	395
2620	GMFCS grade (among infants with cerebral palsy)	 1: I 24% 2: II 23% 3: III 11% 4: IV 14% 5: V 27%	201
2630	Type of cerebral palsy (among infants with cerebral palsy)	 1:Spastic 80% 2:Athetoid 6% 3:Mixed 5% 4:Flaccid 6% 5:Others 3%	254
2640	Topographical distribution (among infants with cerebral palsy)	 1:Quadriplegia 41% 2:Diplegia 28% 3:Paraplegia 18% 4:Hemiplegia 12% 5:Monoplegia 2%	227






No.	Resources of participating hospitals	All hospitals	n
2650	Cause of cerebral palsy (among infants with cerebral palsy)	 4680 663 Others	248
2660	DQ or IQ measurement (among infants with followup at 6 years of age)	 1:Yes 88% 2:No 12%	5343
2670	Reason not to measure DQ or IQ (among infants with followup at 6 years of age)	 1:Normal 44% 2:Developmental delay 25% 3:Refuse by parents 3% 4:Unable to perform 5% 5:Failed to finish 4% 6:Others 19%	519
2680	Method for IQ or DQ measurement (among infants with followup at 6 years of age)	 1:WISCIV 73% 2:WPPSI 2% 3:Tanaka-Binet 6% 4:New Kyoto scale 10% 5:K-ABC 0% 6:Tsumori-Inage 0% 7:Others 7% 8:WISCV 0%	4120
2700	WISCIV Full IQ (mean) (among infants with IQ by WISCIV)	91.6	3005
	SD	14.2	
	95% confidence interval	91.0-92.1	
2702	WISCIV VCI (Verbal) (mean) (among infants with IQ by WISCIV)	94.8	3007
	SD	13.5	
	95% confidence interval	94.3-95.3	

No.	Resources of participating hospitals	All hospitals	n
2704	WISCIV PRI (Perceptual) (mean) (among infants with IQ by WISCIV)	94.6	3016
	SD	14.6	
	95% confidence interval	94.1-95.2	
2706	WISCIV WMI (Working) (mean) (among infants with IQ by WISCIV)	88.9	3011
	SD	16.6	
	95% confidence interval	88.3-89.5	
2708	WISCIV PSI (Processing) (mean) (among infants with IQ by WISCIV)	91.7	3005
	SD	14.6	
	95% confidence interval	91.2-92.2	
2710	WPPSI Full IQ (mean) (among infants with IQ by WIPPSI)	74.0	99
	SD	17.6	
	95% confidence interval	70.6-77.5	
2712	WPPSI VCI (Verbal) (mean) (among infants with IQ by WIPPSI)	74.7	96
	SD	17.0	
	95% confidence interval	71.3-78.1	
2714	WPPSI PRI (Perceptual) (mean) (among infants with IQ by WIPPSI)	81.1	94
	SD	17.2	
	95% confidence interval	77.6-84.5	

No.	Resources of participating hospitals	All hospitals	n
2716	WPPSI PSI (Processing) (mean) (among infants with IQ by WIPPSI)	76.3	84
	SD	19.1	
	95% confidence interval	72.2-80.4	
2718	WPPSI GLC (Global language) (mean) (among infants with IQ by WIPPSI)	80.3	53
	SD	17.8	
	95% confidence interval	75.5-85.0	
2720	WISCV Full scale IQ (mean) (among infants with IQ by WISCV)	0.0	0
	SD	0.0	
	95% confidence interval	0.0-0.0	
2722	WISCV VCI (Verbal) (mean) (among infants with IQ by WISCV)	0.0	0
	SD	0.0	
	95% confidence interval	0.0-0.0	
2724	WISCV VSI (Visual) (mean) (among infants with IQ by WISCV)	0.0	0
	SD	0.0	
	95% confidence interval	0.0-0.0	
2726	WISCV FRI (Fluid) (mean) (among infants with IQ by WISCV)	0.0	0
	SD	0.0	
	95% confidence interval	0.0-0.0	

No.	Resources of participating hospitals	All hospitals	n
2728	WISCV WMI (Working) (mean) (among infants with IQ by WISCV)	0.0	0
	SD	0.0	
	95% confidence interval	0.0-0.0	
2730	WISCV PSI (Processing) (mean) (among infants with IQ by WISCV)	0.0	0
	SD	0.0	
	95% confidence interval	0.0-0.0	
2732	WISCV QRI (Quantitative) (mean) (among infants with IQ by WISCV)	0.0	0
	SD	0.0	
	95% confidence interval	0.0-0.0	
2734	WISCV AWMI (Auditory) (mean) (among infants with IQ by WISCV)	0.0	0
	SD	0.0	
	95% confidence interval	0.0-0.0	
2736	WISCV NVI (Nonverbal) (mean) (among infants with IQ by WISCV)	0.0	0
	SD	0.0	
	95% confidence interval	0.0-0.0	
2738	WISCV GAI (General) (mean) (among infants with IQ by WISCV)	0.0	0
	SD	0.0	
	95% confidence interval	0.0-0.0	

No.	Resources of participating hospitals	All hospitals	n
2740	WISCV CPI (Cognitive) (mean) (among infants with IQ by WISCV)	0.0	0
	SD	0.0	
	95% confidence interval	0.0-0.0	
2820	New Kyoto scale (mean) (among infants with DQ by K scale)	64.6	404
	SD	23.5	
	95% confidence interval	62.4-66.9	
2830	Tanaka-Binet scale (mean) (among infants with IQ by T-B)	74.5	256
	SD	23.1	
	95% confidence interval	71.7-77.3	
2840	K-ABCII scale (mean) (among infants with IQ by K-ABC)	0.0	0
	SD	0.0	
	95% confidence interval	0.0-0.0	
2850	Other method (among infants with IQ or DQ by other methods)	 1:Normal 71% 2:Boderline 20% 3:Delay 9%	161
2860	Evaluated by physician (among infants with followup at 6 years of age)	 1:Normal 61% 2:Boderline 22% 3:Delay 17%	2550

No.	Resources of participating hospitals	All hospitals	n
2870	Asthma (among infants with followup at 6 years of age)	 1: Yes 9% 2: No 91%	4882
2880	Epilepsy (among infants with followup at 6 years of age)	 1: Yes 3% 2: No 97%	4954
2890	Behavioral disorder (among infants with followup at 6 years of age)	 1: Yes 16% 2: No 84%	4876
2892	Type of behavioral disorder (among infants with behavior disorder)	 ADHD 1472 ADHD... 3464 ASD ASD... Others	790
2900	Rehabilitation (among infants with followup at 6 years of age)	 1: Yes 30% 2: No 70%	4936