










No.	Resources of participating hospitals	All hospitals	n
<b>P</b>			
2010	Followup at 1.5 years of age (among infants with alive at discharge)	 <p>1:Yes 29% 2:No 71%</p>	46840
2012	Dead after discharge (among infants with alive at discharge)	 <p>1:Yes 1% 2:No 91% 3:not available 8%</p>	14112
2016	Reason for dropout (among infants with alive at discharge)	 <p>1:Followed at different hospital 84% 2:Admitted in rehabilitation center 0% 3:No contact 15% 4:Others 0% 5:Admitted in other rehabilitation center 0% 6:Hospitalization 0%</p>	2815
2020	Age at followup (mean) (among infants with followup at 1.5 years of age)	1.7	13057
	SD	0.1	
	95% confidence interval	1.7-1.7	
2022	Age corrected at followup (mean) (among infants with followup at 1.5 years of age)	1.5	13283
	SD	0.1	
	95% confidence interval	1.5-1.5	
2030	Body weight (mean) (among infants with followup at 1.5 years of age)	9.3	12735
	SD	1.8	
	95% confidence interval	9.3-9.4	

No.	Resources of participating hospitals	All hospitals	n
2040	Height (mean) (among infants with followup at 1.5 years of age)	77.5	12628
	SD	3.9	
	95% confidence interval	77.4-77.5	
2050	Head circumference (mean) (among infants with followup at 1.5 years of age)	46.4	11712
	SD	1.9	
	95% confidence interval	46.4-46.4	
2060	Oxygen (among infants with followup at 1.5 years of age)	 <p>1:Yes 3% 2:No 97%</p>	12856
2070	Visual impairment (among infants with followup at 1.5 years of age)	 <p>1:Yes 3% 2:No 97%</p>	12606
2080	Cerebral palsy (among infants with followup at 1.5 years of age)	 <p>1:Yes 7% 2:No 93%</p>	12711
2085	DQ measurement (among infants with followup at 1.5 years of age)	 <p>1:Yes 75% 2:No 25%</p>	12704

No.	Resources of participating hospitals	All hospitals	n
2088	Reason not to measure DQ (among infants with DQ measurement)	 <p>1:Normal development by physician diagnosis 58%                  2:Severely damaged by physician diagnosis 16%                  3:Refusal from patents 2%                  4:Impossible to perform due to severely damaged 4%                  5:Failed to perform 4%                  6:others 17%</p>	2849
2100	Method for DQ measurement (among infants with followup at 1.5 years of age)	 <p>1:Kyoto scale 80%                  2:Others 20%</p>	10365
2101	DQ (K scale) (mean) (among infants with DQ measured by K scale)	75.9	6178
	SD	18.0	
	95% confidence interval	75.5-76.4	
2102	DQ corrected age (K scale) (mean) (among infants with DQ measured by K scale)	85.2	7384
	SD	24.0	
	95% confidence interval	84.7-85.8	
2112	DQ (other than K scale) (mean) (among infants with DQ measured by other than K scale)	86.6	1597
	SD	195.4	
	95% confidence interval	77.0-96.2	
2113	DQ corrected age (other than K scale) (mean) (among infants with DQ measured by other than K scale)	88.9	1749
	SD	29.7	
	95% confidence interval	87.6-90.3	