




















No.	Resources of participating hospitals	All hospitals	n
P			
2010	Followup at 1.5 years of age (among infants with alive at discharge)	 <p>1:Yes 28% 2:No 72%</p>	51091
2012	Dead after discharge (among infants with alive at discharge)	 <p>1:Yes 1% 2:No 91% 3:not available 8%</p>	14114
2016	Reason for dropout (among infants with alive at discharge)	 <p>1:Followed at different hospital 84% 2:Admitted in rehabilitation center 0% 3:No contact 15% 4:Others 0% 5:Admitted in other rehabilitation center 0% 6:Hospitalization 0%</p>	2925
2020	Age at followup (mean) (among infants with followup at 1.5 years of age)	1.7	13721
	SD	0.1	
	95% confidence interval	1.7-1.7	
2022	Age corrected at followup (mean) (among infants with followup at 1.5 years of age)	1.5	13947
	SD	0.1	
	95% confidence interval	1.5-1.5	
2030	Body weight (mean) (among infants with followup at 1.5 years of age)	9.3	13385
	SD	2.0	
	95% confidence interval	9.3-9.4	







No.	Resources of participating hospitals	All hospitals	n
2040	Height (mean) (among infants with followup at 1.5 years of age)	77.5	13274
	SD	3.9	
	95% confidence interval	77.4-77.5	
2050	Head circumference (mean) (among infants with followup at 1.5 years of age)	46.4	12314
	SD	1.9	
	95% confidence interval	46.4-46.4	
2060	Oxygen (among infants with followup at 1.5 years of age)	 <p>1:Yes 3% 2:No 97%</p>	13479
2061	Duration of home oxygen (mean) (among infants with oxygen)	14.2	43
	SD	4.3	
	95% confidence interval	12.9-15.5	
2070	Visual impairment (among infants with followup at 1.5 years of age)	 <p>1:Yes 3% 2:No 97%</p>	13218
2071	Severity of visual impairment (among infants with visual impairment)	 <p>1:Less than light perception 12% 2:Amblyopia or ny Stagmus 18% 3:strabismus 54% 4:Others 16%</p>	67



No.	Resources of participating hospitals	All hospitals	n
2072	Eye glasses (among infants with followup at 1.5 years of age)	 <p>1:Yes 2%</p> <p>2:No 98%</p>	1631
2080	Cerebral palsy (among infants with followup at 1.5 years of age)	 <p>1:Yes 7%</p> <p>2:No 93%</p>	13323
2081	GMFCS grade (among infants with cerebral palsy)	 <p>1:I 15%</p> <p>2: 11%</p> <p>3: 9%</p> <p>4: 19%</p> <p>5: 46%</p>	80
2082	Type of cerebral palsy (among infants with cerebral palsy)	 <p>1:Spastic 75%</p> <p>2:Athetoid 8%</p> <p>3:Mixed 4%</p> <p>4:Flaccid 10%</p> <p>5:others 3%</p>	106
2083	Cause of cerebral palsy (among infants with cerebral palsy)	 <p>1:PVL 47%</p> <p>2:IVH 30%</p> <p>3:Others 23%</p>	92
2085	DQ measurement (among infants with followup at 1.5 years of age)	 <p>1:Yes 76%</p> <p>2:No 24%</p>	13373

No.	Resources of participating hospitals	All hospitals	n
2088	Reason not to measure DQ (among infants with DQ measurement)	 <p>1:Normal development by physician diagnosis 58% 2:Severely damaged by physician diagnosis 15% 3:Refusal from patents 2% 4:Impossible to perform due to severely damaged 4% 5:Failed to perform 4% 6:others 17%</p>	2924
2100	Method for DQ measurement (among infants with followup at 1.5 years of age)	 <p>1:Kyoto scale 80% 2:Others 20%</p>	10941
2101	DQ (K scale)(mean) (among infants with DQ measured by K scale)	76.2	6605
	SD	17.7	
	95% confidence interval	75.7-76.6	
2102	DQ corrected age (K scale)(mean) (among infants with DQ measured by K scale)	85.7	7867
	SD	26.5	
	95% confidence interval	85.1-86.3	
2103	DQ postural-motor (K scale)(mean) (among infants with DQ measured by K scale)	75.7	1092
	SD	17.2	
	95% confidence interval	74.7-76.7	
2104	DQ postural-motor corrected age (K scale) (mean) (among infants with DQ measured by K scale)	85.9	1299
	SD	19.1	
	95% confidence interval	84.8-86.9	

No.	Resources of participating hospitals	All hospitals	n
2105	DQ cognitive-adaptive (K scale) (mean) (among infants with DQ measured by K scale)	78.7	1092
	SD	15.2	
	95% confidence interval	77.8-79.6	
2106	DQ cognitive-adaptive corrected age (K scale) (mean) (among infants with DQ measured by K scale)	88.9	1297
	SD	16.1	
	95% confidence interval	88.0-89.8	
2107	DQ language-social (K scale) (mean) (among infants with DQ measured by K scale)	76.4	1093
	SD	16.5	
	95% confidence interval	75.4-77.4	
2108	DQ language-social corrected age (K scale) (mean) (among infants with DQ measured by K scale)	86.4	1299
	SD	17.8	
	95% confidence interval	85.4-87.3	
2111	Method for DQ measurement other than K scale (among infants with DQ measured by other than K scale)	<p>1: Bayley 2% 2: Enjogi 50% 3: Tsumori-Image 24% 4: Others 24%</p>	1970
2112	DQ (other than K scale) (mean) (among infants with DQ measured by other than K scale)	87.5	1655
	SD	191.8	
	95% confidence interval	78.2-96.7	

No.	Resources of participating hospitals	All hospitals	n
2113	DQ corrected age (other than K scale) (mean) (among infants with DQ measured by other than K scale)	90.4	1775
	SD	27.9	
	95% confidence interval	89.1-91.7	
2114	Evaluation (other than K scale) (among infants with DQ measured by other than K scale)	 <p>1:Normal 65% 2:Borderline 14% 3:Delayed 21%</p>	280
2115	Evaluation by physician (among infants with DQ measured by other than K scale)	 <p>1:Normal 65% 2:Borderline 14% 3:Delayed 21%</p>	274
2120	Hearing impairment (among infants with followup at 1.5 years of age)	 <p>1:Yes 2% 2:No 98%</p>	1747
2122	Hearing aide (among infants with hearing impairment)	 <p>1:Yes 25% 2:No 75%</p>	24
2123	Audiometry (among infants with hearing impairment)	 <p>1:Normal 39% 2:Moderate 56% 3:Severe 6%</p>	18

No.	Resources of participating hospitals	All hospitals	n
2130	Asthme (among infants with followup at 1.5 years of age)	 <p>1:Yes 8% 2:No 92%</p>	1776
2140	Epilepsy (among infants with followup at 1.5 years of age)	 <p>1:Yes 2% 2:No 98%</p>	1866
2150	Home medical care (among infants with followup at 1.5 years of age)	 <p>1:Yes 5% 2:No 95%</p>	1841
2151	Mechanical ventilation (among infants with home medical care)	 <p>1:Yes 4% 2:No 96%</p>	54
2152	Tracheostomy (among infants with home medical care)	 <p>1:Yes 5% 2:No 95%</p>	61
2153	Tube feeding (among infants with home medical care)	 <p>1:Yes 21% 2:No 79%</p>	39

No.	Resources of participating hospitals	All hospitals	n
2154	VP shunt (among infants with home medical care)	 <p>1: Yes 9%</p> <p>2: No 91%</p>	55
2160	Rehabilitation (among infants with followup at 1.5 years of age)	 <p>1: Yes 13%</p> <p>2: No 87%</p>	1764