





















No.	Resources of participating hospitals	All hospitals	n
P			
2010	Followup at 1.5 years of age (among infants with alive at discharge)	 <p>1:Yes 35% 2:No 65%</p>	65181
2012	Dead after discharge (among infants with alive at discharge)	 <p>1:Yes 2% 2:No 98% 3:not available 0%</p>	14315
2016	Reason for dropout (among infants with alive at discharge)	 <p>1:Followed at different hospital 74% 2:Admitted in rehabilitation center 0% 3:No contact 13% 4:Others 12% 5:Admitted in other rehabilitation center 0% 6:Hospitalization 0%</p>	5250
2020	Age at followup (mean) (among infants with followup at 1.5 years of age)	1.7	22092
	SD	0.1	
	95% confidence interval	1.7-1.7	
2022	Age corrected at followup (mean) (among infants with followup at 1.5 years of age)	1.5	22337
	SD	0.1	
	95% confidence interval	1.5-1.5	
2030	Body weight (mean) (among infants with followup at 1.5 years of age)	9.4	21713
	SD	1.3	
	95% confidence interval	9.3-9.4	







No.	Resources of participating hospitals	All hospitals	n
2040	Height (mean) (among infants with followup at 1.5 years of age)	77.7	21555
	SD	3.9	
	95% confidence interval	77.6-77.7	
2050	Head circumference (mean) (among infants with followup at 1.5 years of age)	46.4	20043
	SD	1.9	
	95% confidence interval	46.4-46.5	
2060	Oxygen (among infants with followup at 1.5 years of age)	 <p>1:Yes 3% 2:No 97%</p>	21671
2061	Duration of home oxygen (mean) (among infants with oxygen)	14.1	193
	SD	5.1	
	95% confidence interval	13.3-14.8	
2070	Visual impairment (among infants with followup at 1.5 years of age)	 <p>1:Yes 4% 2:No 96%</p>	21321
2071	Severety of visual impairment (among infants with visual impairment)	 <p>1:Less than light perception 7% 2:Amblyopia or ny stagmus 24% 3:strabismus 43% 4:Others 25%</p>	341



No.	Resources of participating hospitals	All hospitals	n
2072	Eye glasses (among infants with followup at 1.5 years of age)	 <p>1:Yes 2% 2:No 98%</p>	9644
2080	Cerebral palsy (among infants with followup at 1.5 years of age)	 <p>1:Yes 6% 2:No 94%</p>	21644
2081	GMFCS grade (among infants with cerebral palsy)	 <p>1:I 37% 2: II 42% 3:III 20% 4:IV 0% 5:V 0%</p>	139
2082	Type of cerebral palsy (among infants with cerebral palsy)	 <p>1:Spastic 75% 2:Athetoid 4% 3:Mixed 5% 4:Flaccid 7% 5:others 9%</p>	510
2083	Cause of cerebral palsy (among infants with cerebral palsy)	 <p>1:PVL 48% 2:IVH 26% 3:Others 25%</p>	493
2085	DQ measurement (among infants with followup at 1.5 years of age)	 <p>1:Yes 81% 2:No 19%</p>	21806

No.	Resources of participating hospitals	All hospitals	n
2088	Reason not to measure DQ (among infants with DQ measurement)	 <p>1:Normal development by physician diagnosis 57% 2:Severely damaged by physician diagnosis 16% 3:Refusal from parents 2% 4:Impossible to perform due to severely damaged 4% 5:Failed to perform 4% 6:others 17%</p>	3760
2100	Method for DQ measurement (among infants with followup at 1.5 years of age)	 <p>1:Kyoto scale 83% 2:Others 17%</p>	18409
2101	DQ (K scale) (mean) (among infants with DQ measured by K scale)	78.5	11075
	SD	14.6	
	95% confidence interval	78.2-78.8	
2102	DQ corrected age (K scale) (mean) (among infants with DQ measured by K scale)	88.4	14011
	SD	15.6	
	95% confidence interval	88.1-88.7	
2103	DQ postural-motor (K scale) (mean) (among infants with DQ measured by K scale)	0.0	0
	SD	0.0	
	95% confidence interval	0.0-0.0	
2104	DQ postural-motor corrected age (K scale) (mean) (among infants with DQ measured by K scale)	86.5	8153
	SD	18.5	
	95% confidence interval	86.1-86.9	

No.	Resources of participating hospitals	All hospitals	n
2105	DQ cognitive-adaptive (K scale) (mean) (among infants with DQ measured by K scale)	80.4	6209
	SD	15.6	
	95% confidence interval	80.0-80.8	
2106	DQ cognitive-adaptive corrected age (K scale) (mean) (among infants with DQ measured by K scale)	90.3	8185
	SD	16.6	
	95% confidence interval	89.9-90.6	
2107	DQ language-social (K scale) (mean) (among infants with DQ measured by K scale)	77.4	6213
	SD	16.4	
	95% confidence interval	77.0-77.8	
2108	DQ language-social corrected age (K scale) (mean) (among infants with DQ measured by K scale)	86.8	8182
	SD	17.5	
	95% confidence interval	86.5-87.2	
2111	Method for DQ measurement other than K scale (among infants with DQ measured by other than K scale)	 <p>1: Bayley 5% 2: Enjogi 53% 3: Tsumori-Inage 26% 4: Others 17%</p>	2995
2112	DQ (other than K scale) (mean) (among infants with DQ measured by other than K scale)	84.5	2238
	SD	19.6	
	95% confidence interval	83.7-85.4	

No.	Resources of participating hospitals	All hospitals	n
2113	DQ corrected age (other than K scale) (mean) (among infants with DQ measured by other than K scale)	94.5	2359
	SD	20.3	
	95% confidence interval	93.6-95.3	
2114	Evaluation (other than K scale) (among infants with DQ measured by other than K scale)	 <p>1:Normal 68% 2:Bordeline 17% 3:Delayed 14%</p>	1230
2115	Evaluation by physician (among infants with DQ measured by other than K scale)	 <p>1:Normal 71% 2:Bordeline 15% 3:Delayed 14%</p>	1087
2120	Hearing impairment (among infants with followup at 1.5 years of age)	 <p>1:Yes 2% 2:No 98%</p>	10253
2122	Hearing aid (among infants with hearing impairment)	 <p>1:Yes 33% 2:No 67%</p>	168
2123	Audiometry (among infants with hearing impairment)	 <p>1:Normal 12% 2:Moderate 69% 3:Severe 19%</p>	141

No.	Resources of participating hospitals	All hospitals		n
2130	Asthme (among infants with followup at 1.5 years of age)		1:Yes 7% 2:No 93%	10184
2140	Epilepsy (among infants with followup at 1.5 years of age)		1:Yes 2% 2:No 98%	9073
2150	Home medical care (among infants with followup at 1.5 years of age)		1:Yes 5% 2:No 95%	8971
2151	Mechanical ventilation (among infants with home medical care)		1:Yes 7% 2:No 93%	313
2152	Tracheostomy (among infants with home medical care)		1:Yes 12% 2:No 88%	311
2153	Tube feeding (among infants with home medical care)		1:Yes 27% 2:No 73%	246

No.	Resources of participating hospitals	All hospitals	n
2154	VP shunt (among infants with home medical care)	 <p>1:Yes 12% 2:No 88%</p>	303
2160	Rehabilitation (among infants with followup at 1.5 years of age)	 <p>1:Yes 14% 2:No 86%</p>	8808