




















No.	Resources of participating hospitals	All hospitals	n
<b>P</b>			
<b>2010</b>	Followup at 1.5 years of age (among infants with alive at discharge)	 <p>1:Yes 34% 2:No 66%</p>	72928
<b>2012</b>	Dead after discharge (among infants with alive at discharge)	 <p>1:Yes 2% 2:No 98% 3:not available 0%</p>	14630
<b>2016</b>	Reason for dropout (among infants with alive at discharge)	 <p>1:Followed at different hospital 74% 2:Admitted in rehabilitation center 0% 3:No contact 13% 4:Others 12% 5:Admitted in other rehabilitation center 0% 6:Hospitalization 0%</p>	5706
<b>2020</b>	Age at followup (mean) (among infants with followup at 1.5 years of age)	1.7	24533
	SD	0.1	
	95% confidence interval	1.7-1.7	
<b>2022</b>	Age corrected at followup (mean) (among infants with followup at 1.5 years of age)	1.5	24774
	SD	0.1	
	95% confidence interval	1.5-1.5	
<b>2030</b>	Body weight (mean) (among infants with followup at 1.5 years of age)	9.4	24113
	SD	1.3	
	95% confidence interval	9.4-9.4	







No.	Resources of participating hospitals	All hospitals	n
2040	Height (mean) (among infants with followup at 1.5 years of age)	77.7	23949
	SD	3.9	
	95% confidence interval	77.7-77.8	
2050	Head circumference (mean) (among infants with followup at 1.5 years of age)	46.5	22170
	SD	1.9	
	95% confidence interval	46.4-46.5	
2060	Oxygen (among infants with followup at 1.5 years of age)	 <p>1: Yes 3% 2: No 97%</p>	23973
2061	Duration of home oxygen (mean) (among infants with oxygen)	14.1	230
	SD	5.2	
	95% confidence interval	13.4-14.7	
2070	Visual impairment (among infants with followup at 1.5 years of age)	 <p>1: Yes 4% 2: No 96%</p>	23595
2071	Severity of visual impairment (among infants with visual impairment)	 <p>1: Less than light perception 7% 2: Amblyopia or nystagmus 24% 3: strabismus 42% 4: Others 27%</p>	415



No.	Resources of participating hospitals	All hospitals	n
2072	Eye glasses (among infants with followup at 1.5 years of age)	 <p>1: Yes 2%</p> <p>2: No 98%</p>	11811
2080	Cerebral palsy (among infants with followup at 1.5 years of age)	 <p>1: Yes 6%</p> <p>2: No 94%</p>	23959
2081	GMFCS grade (among infants with cerebral palsy)	 <p>1: I 38%</p> <p>2: II 41%</p> <p>3: III 21%</p> <p>4: IV 0%</p> <p>5: V 0%</p>	160
2082	Type of cerebral palsy (among infants with cerebral palsy)	 <p>1: Spastic 76%</p> <p>2: Athetoid 4%</p> <p>3: Mixed 5%</p> <p>4: Flaccid 7%</p> <p>5: others 8%</p>	588
2083	Cause of cerebral palsy (among infants with cerebral palsy)	 <p>1: PVL 49%</p> <p>2: IVH 25%</p> <p>3: Others 26%</p>	563
2085	DQ measurement (among infants with followup at 1.5 years of age)	 <p>1: Yes 82%</p> <p>2: No 18%</p>	24244

No.	Resources of participating hospitals	All hospitals	n
2088	Reason not to measure DQ (among infants with DQ measurement)	 <p>1:Normal development by physician diagnosis 58%                  2:Severely damaged by physician diagnosis 16%                  3:Refusal from parents 2%                  4:Impossible to perform due to severely damaged 4%                  5:Failed to perform 4%                  6:others 16%</p>	4019
2100	Method for DQ measurement (among infants with followup at 1.5 years of age)	 <p>1:Kyoto scale 83%                  2:Others 17%</p>	20546
2101	DQ (K scale) (mean) (among infants with DQ measured by K scale)	78.7	12689
	SD	14.6	
	95% confidence interval	78.4-79.0	
2102	DQ corrected age (K scale) (mean) (among infants with DQ measured by K scale)	88.5	15789
	SD	15.5	
	95% confidence interval	88.3-88.8	
2103	DQ postural-motor (K scale) (mean) (among infants with DQ measured by K scale)	0.0	0
	SD	0.0	
	95% confidence interval	0.0-0.0	
2104	DQ postural-motor corrected age (K scale) (mean) (among infants with DQ measured by K scale)	86.8	9947
	SD	18.3	
	95% confidence interval	86.4-87.2	

No.	Resources of participating hospitals	All hospitals	n
2105	DQ cognitive-adaptive (K scale) (mean) (among infants with DQ measured by K scale)	80.6	7841
	SD	15.6	
	95% confidence interval	80.2-80.9	
2106	DQ cognitive-adaptive corrected age (K scale) (mean) (among infants with DQ measured by K scale)	90.4	9976
	SD	16.6	
	95% confidence interval	90.0-90.7	
2107	DQ language-social (K scale) (mean) (among infants with DQ measured by K scale)	77.4	7844
	SD	16.3	
	95% confidence interval	77.0-77.7	
2108	DQ language-social corrected age (K scale) (mean) (among infants with DQ measured by K scale)	86.8	9976
	SD	17.3	
	95% confidence interval	86.5-87.1	
2111	Method for DQ measurement other than K scale (among infants with DQ measured by other than K scale)	<p>1: Bayley 7% 2: Enjogi 52% 3: Tsumori-Inage 26% 4: Others 16%</p>	3269
2112	DQ (other than K scale) (mean) (among infants with DQ measured by other than K scale)	84.4	2403
	SD	19.4	
	95% confidence interval	83.6-85.2	

No.	Resources of participating hospitals	All hospitals	n
2113	DQ corrected age (other than K scale) (mean) (among infants with DQ measured by other than K scale)	94.2	2521
	SD	20.2	
	95% confidence interval	93.4-95.0	
2114	Evaluation (other than K scale) (among infants with DQ measured by other than K scale)	 <p>1:Normal 67% 2:Borderline 18% 3:Delayed 15%</p>	1486
2115	Evaluation by physician (among infants with DQ measured by other than K scale)	 <p>1:Normal 71% 2:Borderline 15% 3:Delayed 14%</p>	1138
2120	Hearing impairment (among infants with followup at 1.5 years of age)	 <p>1:Yes 2% 2:No 98%</p>	12528
2122	Hearing aid (among infants with hearing impairment)	 <p>1:Yes 34% 2:No 66%</p>	211
2123	Audiometry (among infants with hearing impairment)	 <p>1:Normal 16% 2:Moderate 63% 3:Severe 20%</p>	176

No.	Resources of participating hospitals	All hospitals	n
2130	Asthme (among infants with followup at 1.5 years of age)	 <p>1: Yes 6% 2: No 94%</p>	12490
2140	Epilepsy (among infants with followup at 1.5 years of age)	 <p>1: Yes 2% 2: No 98%</p>	11410
2150	Home medical care (among infants with followup at 1.5 years of age)	 <p>1: Yes 4% 2: No 96%</p>	11275
2151	Mechanical ventilation (among infants with home medical care)	 <p>1: Yes 7% 2: No 93%</p>	362
2152	Tracheostomy (among infants with home medical care)	 <p>1: Yes 11% 2: No 89%</p>	359
2153	Tube feeding (among infants with home medical care)	 <p>1: Yes 27% 2: No 73%</p>	278

No.	Resources of participating hospitals	All hospitals	n
2154	VP shunt (among infants with home medical care)	 <p>1: Yes 10%</p> <p>2: No 90%</p>	367
2160	Rehabilitation (among infants with followup at 1.5 years of age)	 <p>1: Yes 14%</p> <p>2: No 86%</p>	11092