















No.	Resources of participating hospitals	All hospitals	n
P			
2010	Followup at 1.5 years of age (among infants with alive at discharge)	 <p>1:Yes 34% 2:No 66%</p>	79642
2012	Dead after discharge (among infants with alive at discharge)	 <p>1:Yes 3% 2:No 97% 3:not available 0%</p>	14780
2016	Reason for dropout (among infants with alive at discharge)	 <p>1:Followed at different hospital 74% 2:Admitted in rehabilitation center 0% 3:No contact 14% 4:Others 12% 5:Admitted in other rehabilitation center 0% 6:Hospitalization 0%</p>	5989
2020	Age at followup (mean) (among infants with followup at 1.5 years of age)	1.7	26214
	SD	0.1	
	95% confidence interval	1.7-1.7	
2022	Age corrected at followup (mean) (among infants with followup at 1.5 years of age)	1.5	26454
	SD	0.1	
	95% confidence interval	1.5-1.5	
2030	Body weight (mean) (among infants with followup at 1.5 years of age)	9.4	25770
	SD	1.3	
	95% confidence interval	9.4-9.4	

No.	Resources of participating hospitals	All hospitals	n
2040	Height (mean) (among infants with followup at 1.5 years of age)	77.8	25600
	SD	3.9	
	95% confidence interval	77.7-77.8	
2050	Head circumference (mean) (among infants with followup at 1.5 years of age)	46.5	23690
	SD	1.9	
	95% confidence interval	46.4-46.5	
2060	Oxygen (among infants with followup at 1.5 years of age)	 <p>1: Yes 3% 2: No 97%</p>	25548
2061	Duration of home oxygen (mean) (among infants with oxygen)	14.0	267
	SD	5.2	
	95% confidence interval	13.4-14.6	
2070	Visual impairment (among infants with followup at 1.5 years of age)	 <p>1: Yes 4% 2: No 96%</p>	25146
2071	Severity of visual impairment (among infants with visual impairment)	 <p>1: Less than light perception 6% 2: Amblyopia or nystagmus 22% 3: strabismus 44% 4: Others 27%</p>	463

Analysis results on infants born in 2003-2022


1.5 years of age V-3/8

No.	Resources of participating hospitals	All hospitals	n
2072	Eye glasses (among infants with followup at 1.5 years of age)	 <p>1: Yes 2% 2: No 98%</p>	13312
2080	Cerebral palsy (among infants with followup at 1.5 years of age)	 <p>1: Yes 6% 2: No 94%</p>	25533
2081	GMFCS grade (among infants with cerebral palsy)	 <p>1: I 36% 2: II 41% 3: III 23% 4: IV 0% 5: V 0%</p>	184
2082	Type of cerebral palsy (among infants with cerebral palsy)	 <p>1: Spastic 76% 2: Athetoid 3% 3: Mixed 5% 4: Flaccid 7% 5: others 8%</p>	645
2083	Cause of cerebral palsy (among infants with cerebral palsy)	 <p>1: PVL 50% 2: IVH 25% 3: Others 25%</p>	617
2085	DQ measurement (among infants with followup at 1.5 years of age)	 <p>1: Yes 82% 2: No 18%</p>	25921

No.	Resources of participating hospitals	All hospitals	n
2088	Reason not to measure DQ (among infants with DQ measurement)	 <p>1:Normal development by physician diagnosis 58% 2:Severely damaged by physician diagnosis 16% 3:Refusal from patents 2% 4:Impossible to perform due to severely damaged 4% 5:Failed to perform 4% 6:others 16%</p>	4157
2100	Method for DQ measurement (among infants with followup at 1.5 years of age)	 <p>1:Kyoto scale 83% 2:Others 17%</p>	22047
2101	DQ (K scale) (mean) (among infants with DQ measured by K scale)	78.7	13872
	SD	14.6	
	95% confidence interval	78.5-79.0	
2102	DQ corrected age (K scale) (mean) (among infants with DQ measured by K scale)	88.5	16989
	SD	15.5	
	95% confidence interval	88.3-88.8	
2103	DQ postural-motor (K scale) (mean) (among infants with DQ measured by K scale)	0.0	0
	SD	0.0	
	95% confidence interval	0.0-0.0	
2104	DQ postural-motor corrected age (K scale) (mean) (among infants with DQ measured by K scale)	86.9	11157
	SD	18.3	
	95% confidence interval	86.6-87.3	






Analysis results on infants born in 2003-2022







1.5 years of age V-5/8

No.	Resources of participating hospitals	All hospitals	n
2105	DQ cognitive-adaptive (K scale) (mean) (among infants with DQ measured by K scale)	80.5	9033
	SD	15.6	
	95% confidence interval	80.2-80.9	
2106	DQ cognitive-adaptive corrected age (K scale) (mean) (among infants with DQ measured by K scale)	90.3	11180
	SD	16.6	
	95% confidence interval	90.0-90.6	
2107	DQ language-social (K scale) (mean) (among infants with DQ measured by K scale)	77.3	9037
	SD	16.3	
	95% confidence interval	77.0-77.6	
2108	DQ language-social corrected age (K scale) (mean) (among infants with DQ measured by K scale)	86.7	11185
	SD	17.3	
	95% confidence interval	86.4-87.0	
2111	Method for DQ measurement other than K scale (among infants with DQ measured by other than K scale)	 <p>1: Bayley 7% 2: Enjogi 52% 3: Tsumori-Inage 27% 4: Others 15%</p>	3491
2112	DQ (other than K scale) (mean) (among infants with DQ measured by other than K scale)	84.5	2560
	SD	19.2	
	95% confidence interval	83.8-85.3	

Analysis results on infants born in 2003-2022



1.5 years of age V-6/8

No.	Resources of participating hospitals	All hospitals	n
2113	DQ corrected age (other than K scale) (mean) (among infants with DQ measured by other than K scale)	94.3	2677
	SD	20.0	
	95% confidence interval	93.6-95.1	
2114	Evaluation (other than K scale) (among infants with DQ measured by other than K scale)	 <p>1:Normal 67% 2:Borderline 18% 3:Delayed 14%</p>	1696
2115	Evaluation by physician (among infants with DQ measured by other than K scale)	 <p>1:Normal 71% 2:Borderline 15% 3:Delayed 14%</p>	1137
2120	Hearing impairment (among infants with followup at 1.5 years of age)	 <p>1:Yes 2% 2:No 98%</p>	14069
2122	Hearing aid (among infants with hearing impairment)	 <p>1:Yes 33% 2:No 67%</p>	236
2123	Audiometry (among infants with hearing impairment)	 <p>1:Normal 19% 2:Moderate 62% 3:Severe 19%</p>	193

No.	Resources of participating hospitals	All hospitals	n
2130	Asthme (among infants with followup at 1.5 years of age)	 <p>1: Yes 6% 2: No 94%</p>	14036
2140	Epilepsy (among infants with followup at 1.5 years of age)	 <p>1: Yes 2% 2: No 98%</p>	12981
2150	Home medical care (among infants with followup at 1.5 years of age)	 <p>1: Yes 4% 2: No 96%</p>	12857
2151	Mechanical ventilation (among infants with home medical care)	 <p>1: Yes 7% 2: No 93%</p>	396
2152	Tracheostomy (among infants with home medical care)	 <p>1: Yes 10% 2: No 90%</p>	396
2153	Tube feeding (among infants with home medical care)	 <p>1: Yes 26% 2: No 74%</p>	306

Analysis results on infants born in 2003-2022

1.5 years of age V-8/8

No.	Resources of participating hospitals	All hospitals	n
2154	VP shunt (among infants with home medical care)	 <p>1: Yes 11% 2: No 89%</p>	406
2160	Rehabilitation (among infants with followup at 1.5 years of age)	 <p>1: Yes 14% 2: No 86%</p>	12651