




















No.	Resources of participating hospitals	All hospitals	n
<b>P</b>			
<b>2010</b>	Followup at 1.5 years of age (among infants with alive at discharge)	 <p>1:Yes 35% 2:No 65%</p>	83568
<b>2012</b>	Dead after discharge (among infants with alive at discharge)	 <p>1:Yes 3% 2:No 97% 3:not available 0%</p>	15044
<b>2016</b>	Reason for dropout (among infants with alive at discharge)	 <p>1:Followed at different hospital 74% 2:Admitted in rehabilitation center 0% 3:No contact 14% 4:Others 11% 5:Admitted in other rehabilitation center 0% 6:Hospitalization 0%</p>	6415
<b>2020</b>	Age at followup (mean) (among infants with followup at 1.5 years of age)	1.7	28280
	SD	0.1	
	95% confidence interval	1.7-1.7	
<b>2022</b>	Age corrected at followup (mean) (among infants with followup at 1.5 years of age)	1.5	28505
	SD	0.1	
	95% confidence interval	1.5-1.5	
<b>2030</b>	Body weight (mean) (among infants with followup at 1.5 years of age)	9.4	27712
	SD	1.3	
	95% confidence interval	9.4-9.4	







No.	Resources of participating hospitals	All hospitals	n
2040	Height (mean) (among infants with followup at 1.5 years of age)	77.7	27533
	SD	3.9	
	95% confidence interval	77.7-77.8	
2050	Head circumference (mean) (among infants with followup at 1.5 years of age)	46.4	25377
	SD	1.9	
	95% confidence interval	46.4-46.5	
2060	Oxygen (among infants with followup at 1.5 years of age)	 <p>1:Yes 3% 2:No 97%</p>	27452
2061	Duration of home oxygen (mean) (among infants with oxygen)	14.0	297
	SD	5.2	
	95% confidence interval	13.5-14.6	
2070	Visual impairment (among infants with followup at 1.5 years of age)	 <p>1:Yes 4% 2:No 96%</p>	27010
2071	Severety of visual impairment (among infants with visual impairment)	 <p>1:Less than light perception 6% 2:Amblyopia or ny stagmus 24% 3:strabismus 43% 4:Others 27%</p>	535



No.	Resources of participating hospitals	All hospitals	n
2072	Eye glasses (among infants with followup at 1.5 years of age)	 <p>1:Yes 2%</p> <p>2:No 98%</p>	15053
2080	Cerebral palsy (among infants with followup at 1.5 years of age)	 <p>1:Yes 6%</p> <p>2:No 94%</p>	27412
2081	GMFCS grade (among infants with cerebral palsy)	 <p>1:I 18%</p> <p>2:II 19%</p> <p>3:III 12%</p> <p>4:IV 20%</p> <p>5:V 31%</p>	486
2082	Type of cerebral palsy (among infants with cerebral palsy)	 <p>1:Spastic 75%</p> <p>2:Athetoid 4%</p> <p>3:Mixed 5%</p> <p>4:Flaccid 7%</p> <p>5:others 8%</p>	714
2083	Cause of cerebral palsy (among infants with cerebral palsy)	 <p>1:PVL 51%</p> <p>2:IVH 26%</p> <p>3:Others 24%</p>	692
2085	DQ measurement (among infants with followup at 1.5 years of age)	 <p>1:Yes 82%</p> <p>2:No 18%</p>	27988

No.	Resources of participating hospitals	All hospitals	n
2088	Reason not to measure DQ (among infants with DQ measurement)	 <p>1:Normal development by physician diagnosis 58%  2:Severely damaged by physician diagnosis 16%  3:Refusal from parents 2%  4:Impossible to perform due to severely damaged 4%  5:Failed to perform 4%  6:others 16%</p>	4501
2100	Method for DQ measurement (among infants with followup at 1.5 years of age)	 <p>1:Kyoto scale 83%  2:Others 17%</p>	23716
2101	DQ (K scale) (mean) (among infants with DQ measured by K scale)	78.8	15031
	SD	14.6	
	95% confidence interval	78.6-79.1	
2102	DQ corrected age (K scale) (mean) (among infants with DQ measured by K scale)	88.6	18361
	SD	15.6	
	95% confidence interval	88.3-88.8	
2103	DQ postural-motor (K scale) (mean) (among infants with DQ measured by K scale)	77.9	10221
	SD	17.3	
	95% confidence interval	77.6-78.2	
2104	DQ postural-motor corrected age (K scale) (mean) (among infants with DQ measured by K scale)	87.2	12557
	SD	18.5	
	95% confidence interval	86.8-87.5	

No.	Resources of participating hospitals	All hospitals	n
2105	DQ cognitive-adaptive (K scale) (mean) (among infants with DQ measured by K scale)	80.6	10220
	SD	15.7	
	95% confidence interval	80.3-80.9	
2106	DQ cognitive-adaptive corrected age (K scale) (mean) (among infants with DQ measured by K scale)	90.3	12584
	SD	16.7	
	95% confidence interval	90.0-90.6	
2107	DQ language-social (K scale) (mean) (among infants with DQ measured by K scale)	77.2	10226
	SD	16.3	
	95% confidence interval	76.9-77.5	
2108	DQ language-social corrected age (K scale) (mean) (among infants with DQ measured by K scale)	86.6	12591
	SD	17.3	
	95% confidence interval	86.3-86.9	
2111	Method for DQ measurement other than K scale (among infants with DQ measured by other than K scale)	<p>1: Bayley 7% 2: Enjogi 53% 3: Tsumori-Inage 26% 4: Others 14%</p>	3734
2112	DQ (other than K scale) (mean) (among infants with DQ measured by other than K scale)	84.4	2687
	SD	19.2	
	95% confidence interval	83.6-85.1	

No.	Resources of participating hospitals	All hospitals	n
2113	DQ corrected age (other than K scale) (mean) (among infants with DQ measured by other than K scale)	93.9	2864
	SD	20.4	
	95% confidence interval	93.1-94.6	
2114	Evaluation (other than K scale) (among infants with DQ measured by other than K scale)	 <p>1:Normal 66% 2:Borderline 18% 3:Delayed 16%</p>	1931
2115	Evaluation by physician (among infants with DQ measured by other than K scale)	 <p>1:Normal 71% 2:Borderline 15% 3:Delayed 14%</p>	1137
2120	Hearing impairment (among infants with followup at 1.5 years of age)	 <p>1:Yes 2% 2:No 98%</p>	15991
2122	Hearing aid (among infants with hearing impairment)	 <p>1:Yes 33% 2:No 67%</p>	264
2123	Audiometry (among infants with hearing impairment)	 <p>1:Normal 19% 2:Moderate 61% 3:Severe 20%</p>	214

No.	Resources of participating hospitals	All hospitals	n
2130	Asthme (among infants with followup at 1.5 years of age)	 <p>1:Yes 6% 2:No 94%</p>	15963
2140	Epilepsy (among infants with followup at 1.5 years of age)	 <p>1:Yes 2% 2:No 98%</p>	14909
2150	Home medical care (among infants with followup at 1.5 years of age)	 <p>1:Yes 5% 2:No 95%</p>	14793
2151	Mechanical ventilation (among infants with home medical care)	 <p>1:Yes 7% 2:No 93%</p>	466
2152	Tracheostomy (among infants with home medical care)	 <p>1:Yes 9% 2:No 91%</p>	474
2153	Tube feeding (among infants with home medical care)	 <p>1:Yes 25% 2:No 75%</p>	346

No.	Resources of participating hospitals	All hospitals	n
2154	VP shunt (among infants with home medical care)	 <p>1: Yes 10% 2: No 90%</p>	486
2160	Rehabilitation (among infants with followup at 1.5 years of age)	 <p>1: Yes 15% 2: No 85%</p>	14579