















No.	Resources of participating hospitals	All hospitals	n
P			
2010	Followup at 1.5 years of age (among infants with alive at discharge)	 1:Yes 35% 2:No 65%	86535
2012	Dead after discharge (among infants with alive at discharge)	 1:Yes 3% 2:No 97% 3:not available 0%	15082
2016	Reason for dropout (among infants with alive at discharge)	 1:Followed at different hospital 74% 2:Admitted in rehabilitation center 0% 3:No contact 14% 4:Others 11% 5:Admitted in other rehabilitation center 0% 6:Hospitalization 0%	6683
2020	Age at followup (mean) (among infants with followup at 1.5 years of age)	1.7	29973
	SD	0.1	
	95% confidence interval	1.7-1.7	
2022	Age corrected at followup (mean) (among infants with followup at 1.5 years of age)	1.5	30197
	SD	0.1	
	95% confidence interval	1.5-1.5	
2030	Body weight (mean) (among infants with followup at 1.5 years of age)	9.4	29349
	SD	1.3	
	95% confidence interval	9.4-9.4	







No.	Resources of participating hospitals	All hospitals	n
2040	Height (mean) (among infants with followup at 1.5 years of age)	77.8	29166
	SD	3.9	
	95% confidence interval	77.7-77.8	
2050	Head circumference (mean) (among infants with followup at 1.5 years of age)	46.5	26793
	SD	1.9	
	95% confidence interval	46.4-46.5	
2060	Oxygen (among infants with followup at 1.5 years of age)	 1:Yes 4% 2:No 96%	29018
2061	Duration of home oxygen (mean) (among infants with oxygen)	14.1	328
	SD	5.2	
	95% confidence interval	13.5-14.6	
2070	Visual impairment (among infants with followup at 1.5 years of age)	 1:Yes 4% 2:No 96%	28574
2071	Severity of visual impairment (among infants with visual impairment)	 1:Less than light perception 5% 2:Amblyopia or nystagmus 23% 3:strabismus 42% 4:Others 30%	621



No.	Resources of participating hospitals	All hospitals	n
2072	Eye glasses (among infants with followup at 1.5 years of age)	1: Yes 2% 2: No 98%	16618
2080	Cerebral palsy (among infants with followup at 1.5 years of age)	1: Yes 6% 2: No 94%	28992
2081	GMFCS grade (among infants with cerebral palsy)	1: I 18% 2: II 19% 3: III 12% 4: IV 21% 5: V 30%	524
2082	Type of cerebral palsy (among infants with cerebral palsy)	1: Spastic 76% 2: Athetoid 3% 3: Mixed 5% 4: Flaccid 7% 5: others 8%	760
2083	Cause of cerebral palsy (among infants with cerebral palsy)	1: PVL 51% 2: IVH 26% 3: Others 24%	730
2085	DQ measurement (among infants with followup at 1.5 years of age)	1: Yes 83% 2: No 17%	29676

No.	Resources of participating hospitals	All hospitals	n
2088	Reason not to measure DQ (among infants with DQ measurement)	 1:Normal development by physician diagnosis 59% 2:Severely damaged by physician diagnosis 16% 3:Refusal from parents 2% 4:Impossible to perform due to severely damaged 4% 5:Failed to perform 4% 6:others 15%	4625
2100	Method for DQ measurement (among infants with followup at 1.5 years of age)	 1:Kyoto scale 84% 2:Others 16%	25212
2101	DQ (K scale) (mean) (among infants with DQ measured by K scale)	78.8	16221
	SD	14.7	
	95% confidence interval	78.5-79.0	
2102	DQ corrected age (K scale) (mean) (among infants with DQ measured by K scale)	88.6	19658
	SD	15.6	
	95% confidence interval	88.3-88.8	
2103	DQ postural-motor (K scale) (mean) (among infants with DQ measured by K scale)	77.8	11427
	SD	17.3	
	95% confidence interval	77.5-78.1	
2104	DQ postural-motor corrected age (K scale) (mean) (among infants with DQ measured by K scale)	87.2	13866
	SD	18.6	
	95% confidence interval	86.9-87.5	

No.	Resources of participating hospitals	All hospitals	n
2105	DQ cognitive-adaptive (K scale) (mean) (among infants with DQ measured by K scale)	80.5	11423
	SD	15.7	
	95% confidence interval	80.2-80.8	
2106	DQ cognitive-adaptive corrected age (K scale) (mean) (among infants with DQ measured by K scale)	90.2	13894
	SD	16.8	
	95% confidence interval	89.9-90.5	
2107	DQ language-social (K scale) (mean) (among infants with DQ measured by K scale)	77.0	11431
	SD	16.3	
	95% confidence interval	76.7-77.3	
2108	DQ language-social corrected age (K scale) (mean) (among infants with DQ measured by K scale)	86.5	13903
	SD	17.4	
	95% confidence interval	86.2-86.8	
2111	Method for DQ measurement other than K scale (among infants with DQ measured by other than K scale)	 1: Bayley 8% 2: Enjogi 53% 3: Tsumori-Inage 26% 4: Others 13%	3826
2112	DQ (other than K scale) (mean) (among infants with DQ measured by other than K scale)	84.3	2701
	SD	19.3	
	95% confidence interval	83.6-85.0	

No.	Resources of participating hospitals	All hospitals	n
2113	DQ corrected age (other than K scale) (mean) (among infants with DQ measured by other than K scale)	93.8	2892
	SD	20.5	
	95% confidence interval	93.1-94.6	
2114	Evaluation (other than K scale) (among infants with DQ measured by other than K scale)	 1: Normal 66% 2: Borderline 19% 3: Delayed 16%	2065
2115	Evaluation by physician (among infants with DQ measured by other than K scale)	 1: Normal 71% 2: Borderline 15% 3: Delayed 14%	1134
2120	Hearing impairment (among infants with followup at 1.5 years of age)	 1: Yes 2% 2: No 98%	17647
2122	Hearing aid (among infants with hearing impairment)	 1: Yes 33% 2: No 67%	284
2123	Audiometry (among infants with hearing impairment)	 1: Normal 20% 2: Moderate 60% 3: Severe 20%	225

No.	Resources of participating hospitals	All hospitals	n
2130	Asthme (among infants with followup at 1.5 years of age)	 1:Yes 6% 2:No 94%	17628
2140	Epilepsy (among infants with followup at 1.5 years of age)	 1:Yes 2% 2:No 98%	16595
2150	Home medical care (among infants with followup at 1.5 years of age)	 1:Yes 5% 2:No 95%	16462
2151	Mechanical ventilation (among infants with home medical care)	 1:Yes 6% 2:No 94%	513
2152	Tracheostomy (among infants with home medical care)	 1:Yes 9% 2:No 91%	524
2153	Tube feeding (among infants with home medical care)	 1:Yes 25% 2:No 75%	379

No.	Resources of participating hospitals	All hospitals	n
2154	VP shunt (among infants with home medical care)	 1: Yes 9% 2: No 91%	543
2160	Rehabilitation (among infants with followup at 1.5 years of age)	 1: Yes 15% 2: No 85%	16266