



















No.	Resources of participating hospitals	All hospitals	n
Q			
2210	Followup at 3 years of age (among infants with alive at discharge)	 <p>1:Yes 36% 2:No 64%</p>	51091
2212	Dead after discharge (among infants with alive at discharge)	 <p>1:Yes 1% 2:No 89% 3:not available 10%</p>	16540
2216	Reason for dropout (among infants with alive at discharge)	 <p>1:Followed at different hospital 72% 2:Admitted in rehabilitation center 1% 3:No contact 28% 4:Others 0% 5:Admitted in other rehabilitation center 0% 6:Hospitalization 0%</p>	3819
2220	Age at followup (mean) (among infants with followup at 3 years of age)	3.1	17891
	SD	0.2	
	95% confidence interval	3.1-3.1	
2222	Age corrected at followup (mean) (among infants with followup at 3 years of age)	2.3	1266
	SD	0.5	
	95% confidence interval	2.3-2.4	
2230	Body weight (mean) (among infants with followup at 3 years of age)	12.2	17716
	SD	3.4	
	95% confidence interval	12.2-12.3	







No.	Resources of participating hospitals	All hospitals	n
2240	Height (mean) (among infants with followup at 3 years of age)	89.3	17470
	SD	4.3	
	95% confidence interval	89.2-89.4	
2250	Head circumference (mean) (among infants with followup at 3 years of age)	48.3	15946
	SD	2.0	
	95% confidence interval	48.3-48.3	
2252	Chest circumference (mean) (among infants with followup at 3 years of age)	48.7	9454
	SD	2.7	
	95% confidence interval	48.6-48.7	
2254	Abdominal circumference (mean) (among infants with followup at 3 years of age)	46.2	4164
	SD	3.4	
	95% confidence interval	46.1-46.3	
2260	Oxygen (among infants with followup at 3 years of age)	 <p>1:Yes 2% 2:No 98%</p>	14132
2261	Duration of home oxygen (mean) (among infants with oxygen)	33.9	57
	SD	130.4	
	95% confidence interval	0.1-67.8	







No.	Resources of participating hospitals	All hospitals	n
2270	Visual impairment (among infants with followup at 3 years of age)	 <p>1:Yes 6% 2:No 94%</p>	16856
2271	Severity of visual impairment (among infants with visual impairment)	 <p>1:Less than light perception 4% 2:Amblyopia or ny Stagmus 17% 3:strabismus 44% 4:Others 35%</p>	71
2272	Eye glasses (among infants with visual impairment)	 <p>1:Yes 4% 2:No 96%</p>	1121
2280	Cerebral palsy (among infants with followup at 3 years of age)	 <p>1:Yes 8% 2:No 92%</p>	17518
2281	GMFCS grade (among infants with cerebral palsy)	 <p>1:I 22% 2: 20% 3: 4% 4: 7% 5: 47%</p>	55
2282	Type of cerebral palsy (among infants with cerebral palsy)	 <p>1:Spastic 82% 2:Athetoid 4% 3: Mixed 4% 4:Flaccid 10% 5:others 0%</p>	77

No.	Resources of participating hospitals	All hospitals	n
2283	Cause of cerebral palsy (among infants with cerebral palsy)	 <p>1: PVL 63% 2: IVH 26% 3: Others 11%</p>	65
2285	DQ measurement (among infants with followup at 3 years of age)	 <p>1: Yes 86% 2: No 14%</p>	14576
2288	Reason not to measure DQ (among infants with DQ measurement)	 <p>1: Normal development by physician diagnosis 47% 2: Severely damaged by physician diagnosis 18% 3: Refusal from patients 4% 4: Impossible to perform due to severely damaged 8% 5: Failed to perform 10% 6: others 13%</p>	1748
2300	Method for DQ measurement (among infants with followup at 1.5 years of age)	 <p>1: Kyoto scale 86% 2: Others 14%</p>	15908
2301	DQ (K scale) (mean) (among infants with DQ measured by K scale)	83.6	10850
	SD	16.0	
	95% confidence interval	83.3-83.9	
2302	DQ corrected age (K scale) (mean) (among infants with DQ measured by K scale)	88.4	5836
	SD	18.3	
	95% confidence interval	87.9-88.9	

No.	Resources of participating hospitals	All hospitals	n
2303	DQ postural-motor (K scale) (mean) (among infants with DQ measured by K scale)	86.8	8390
	SD	21.1	
	95% confidence interval	86.4-87.3	
2304	DQ postural-motor corrected age (K scale) (mean) (among infants with DQ measured by K scale)	94.2	5812
	SD	45.2	
	95% confidence interval	93.1-95.4	
2305	DQ cognitive-adaptive (K scale) (mean) (among infants with DQ measured by K scale)	85.7	8456
	SD	92.3	
	95% confidence interval	83.7-87.6	
2306	DQ cognitive-adaptive corrected age (K scale) (mean) (among infants with DQ measured by K scale)	111.7	5844
	SD	1447.9	
	95% confidence interval	74.6-148.8	
2307	DQ language-social (K scale) (mean) (among infants with DQ measured by K scale)	82.7	8440
	SD	18.3	
	95% confidence interval	82.3-83.1	
2308	DQ language-social corrected age (K scale) (mean) (among infants with DQ measured by K scale)	91.2	5825
	SD	53.2	
	95% confidence interval	89.8-92.5	

No.	Resources of participating hospitals	All hospitals	n
2311	Method for DQ other than K scale (among infants with DQ measured by other than K scale)	 <p>1: Bayley 1% 2: Enjogi 63% 3: Tsumori-Inage 21% 4: Others 15%</p>	1510
2312	DQ (other than K scale) (mean) (among infants with DQ measured by other than K scale)	87.7	1430
	SD	24.6	
	95% confidence interval	86.4-88.9	
2313	DQ corrected age (other than K scale) (mean) (among infants with DQ measured by other than K scale)	97.7	610
	SD	23.9	
	95% confidence interval	95.8-99.6	
2314	Evaluation (other than K scale) (among infants with DQ measured by other than K scale)	 <p>1: Normal 66% 2: Bordeline 7% 3: Delayed 27%</p>	115
2315	Evaluation by physician (among infants with DQ measured by other than K scale)	 <p>1: Normal 65% 2: Bordeline 7% 3: Delayed 28%</p>	109
2320	Hearing impairment (among infants with followup at 3 years of age)	 <p>1: Yes 2% 2: No 98%</p>	1162

No.	Resources of participating hospitals	All hospitals	n
2322	Hearing aide (among infants with hearing impairment)	 <p>1:Yes 81% 2:No 19%</p>	16
2323	Audiometry (among infants with hearing impairment)	 <p>1:Normal 40% 2:Moderate 40% 3:Severe 20%</p>	15
2330	Asthme (among infants with followup at 3 years of age)	 <p>1:Yes 10% 2:No 90%</p>	1186
2340	Epilepsy (among infants with followup at 3 years of age)	 <p>1:Yes 2% 2:No 98%</p>	1225
2350	Home medical care (among infants with followup at 3 years of age)	 <p>1:Yes 4% 2:No 96%</p>	1210
2351	Mechanical ventilation (among infants with home medical care)	 <p>1:Yes 21% 2:No 79%</p>	42

No.	Resources of participating hospitals	All hospitals	n
2352	Tracheostomy (among infants with home medical care)	 <p>1:Yes 14% 2:No 86%</p>	42
2353	Tube feeding (among infants with home medical care)	 <p>1:Yes 57% 2:No 43%</p>	42
2354	VP shunt (among infants with home medical care)	 <p>1:Yes 26% 2:No 74%</p>	42
2360	Rehabilitation (among infants with followup at 3 years of age)	 <p>1:Yes 23% 2:No 77%</p>	1191
2370	Behavioral disorder (among infants with followup at 3 years of age)	 <p>1:Yes 11% 2:No 89%</p>	1150
2372	Type of behavioral disorder (among infants with followup at 3 years of age)	 <p>1:ADHD or suspected ADHD 31% 2:ASD or suspected ASD 50% 3:Others 18%</p>	121