



















No.	Resources of participating hospitals	All hospitals	n
Q			
2210	Followup at 3 years of age (among infants with alive at discharge)	 <p>1:Yes 39% 2:No 61%</p>	59723
2212	Dead after discharge (among infants with alive at discharge)	 <p>1:Yes 1% 2:No 87% 3:not available 12%</p>	18013
2216	Reason for dropout (among infants with alive at discharge)	 <p>1:Followed at different hospital 60% 2:Admitted in rehabilitaion center 1% 3:No contact 25% 4:Others 15% 5:Admitted in other rehabilitation center 0% 6:Hospitalization 0%</p>	5768
2220	Age at followup (mean) (among infants with followup at 3 years of age)	3 . 1	22916
	SD	0 . 2	
	95% confidence interval	3 . 1 - 3 . 1	
2222	Age corrected at followup (mean) (among infants with followup at 3 years of age)	2 . 3	6907
	SD	0 . 5	
	95% confidence interval	2 . 3 - 2 . 4	
2230	Body weight (mean) (among infants with followup at 3 years of age)	12 . 1	22648
	SD	1 . 7	
	95% confidence interval	12 . 1 - 12 . 1	







No.	Resources of participating hospitals	All hospitals	n
2240	Height (mean) (among infants with followup at 3 years of age)	89.3	22369
	SD	4.2	
	95% confidence interval	89.3-89.4	
2250	Head circumference (mean) (among infants with followup at 3 years of age)	48.3	20330
	SD	2.0	
	95% confidence interval	48.3-48.3	
2252	Chest circumference (mean) (among infants with followup at 3 years of age)	48.7	11615
	SD	2.7	
	95% confidence interval	48.7-48.8	
2254	Abdominal circumference (mean) (among infants with followup at 3 years of age)	46.2	5316
	SD	3.4	
	95% confidence interval	46.1-46.3	
2260	Oxygen (among infants with followup at 3 years of age)	 <p>1: Yes 2% 2: No 98%</p>	18956
2261	Duration of home oxygen (mean) (among infants with oxygen)	17.8	232
	SD	9.5	
	95% confidence interval	16.5-19.0	







No.	Resources of participating hospitals	All hospitals	n
2270	Visual impairment (among infants with followup at 3 years of age)	 <p>1: Yes 6% 2: No 94%</p>	21754
2271	Severety of visual impairment (among infants with visual impairment)	 <p>1: Less than light perception 3% 2: Amblyopia or n y Stagmus 31% 3: strabismus 30% 4: Others 36%</p>	397
2272	Eye glasses (among infants with visual impairment)	 <p>1: Yes 5% 2: No 95%</p>	6652
2280	Cerebral palsy (among infants with followup at 3 years of age)	 <p>1: Yes 8% 2: No 92%</p>	22541
2281	GMFCS grade (among infants with cerebral palsy)	 <p>1: I 26% 2: 20% 3: 10% 4: 16% 5: 29%</p>	200
2282	Type of cerebral palsy (among infants with cerebral palsy)	 <p>1: Spastic 78% 2: Athetoid 4% 3: Mixed 5% 4: Flaccid 7% 5: others 6%</p>	405

No.	Resources of participating hospitals	All hospitals	n
2283	Cause of cerebral palsy (among infants with cerebral palsy)	 <p>1: PVL 51% 2: IVH 28% 3: Others 21%</p>	355
2285	DQ measurement (among infants with followup at 3 years of age)	 <p>1: Yes 90% 2: No 10%</p>	22373
2288	Reason not to measure DQ (among infants with DQ measurement)	 <p>1: Normal development by physician diagnosis 44% 2: Severely damaged by physician diagnosis 20% 3: Refusal from parents 4% 4: Impossible to perform due to severely damaged 8% 5: Failed to perform 10% 6: others 13%</p>	2054
2300	Method for DQ measurement (among infants with followup at 1.5 years of age)	 <p>1: Kyoto scale 87% 2: Others 13%</p>	20703
2301	DQ (K scale) (mean) (among infants with DQ measured by K scale)	83.8	17568
	SD	16.0	
	95% confidence interval	83.5-84.0	
2302	DQ corrected age (K scale) (mean) (among infants with DQ measured by K scale)	88.7	9432
	SD	17.9	
	95% confidence interval	88.3-89.0	

No.	Resources of participating hospitals	All hospitals	n
2303	DQ postural-motor (K scale) (mean) (among infants with DQ measured by K scale)	86.4	12867
	SD	21.2	
	95% confidence interval	86.0-86.7	
2304	DQ postural-motor corrected age (K scale) (mean) (among infants with DQ measured by K scale)	91.3	9355
	SD	23.1	
	95% confidence interval	90.9-91.8	
2305	DQ cognitive-adaptive (K scale) (mean) (among infants with DQ measured by K scale)	84.5	12979
	SD	16.5	
	95% confidence interval	84.2-84.8	
2306	DQ cognitive-adaptive corrected age (K scale) (mean) (among infants with DQ measured by K scale)	89.7	9416
	SD	18.3	
	95% confidence interval	89.3-90.1	
2307	DQ language-social (K scale) (mean) (among infants with DQ measured by K scale)	82.8	12961
	SD	18.5	
	95% confidence interval	82.4-83.1	
2308	DQ language-social corrected age (K scale) (mean) (among infants with DQ measured by K scale)	87.9	9387
	SD	20.2	
	95% confidence interval	87.5-88.3	

No.	Resources of participating hospitals	All hospitals	n
2311	Method for DQ other than K scale (among infants with DQ measured by other than K scale)	 <p>1: Bayley 4% 2: Enjogi 64% 3: Tsumori-Inage 19% 4: Others 13%</p>	2061
2312	DQ (other than K scale) (mean) (among infants with DQ measured by other than K scale)	88.0	2240
	SD	24.3	
	95% confidence interval	86.9-89.0	
2313	DQ corrected age (other than K scale) (mean) (among infants with DQ measured by other than K scale)	97.8	843
	SD	24.1	
	95% confidence interval	96.1-99.4	
2314	Evaluation (other than K scale) (among infants with DQ measured by other than K scale)	 <p>1: Normal 76% 2: Bordeline 10% 3: Delayed 14%</p>	533
2315	Evaluation by physician (among infants with DQ measured by other than K scale)	 <p>1: Normal 74% 2: Bordeline 9% 3: Delayed 17%</p>	392
2320	Hearing impairment (among infants with followup at 3 years of age)	<p>1</p>  <p>1: Yes 0% 2: No 0%</p>	0

No.	Resources of participating hospitals	All hospitals	n
2322	Hearing aide (among infants with hearing impairment)	 <p>1: Yes 0% 2: No 0%</p>	0
2323	Audiometry (among infants with hearing impairment)	 <p>1: Normal 0% 2: Moderate 0% 3: Severe 0%</p>	0
2330	Asthme (among infants with followup at 3 years of age)	 <p>1: Yes 8% 2: No 92%</p>	6625
2340	Epilepsy (among infants with followup at 3 years of age)	 <p>1: Yes 3% 2: No 97%</p>	6861
2350	Home medical care (among infants with followup at 3 years of age)	 <p>1: Yes 4% 2: No 96%</p>	5397
2351	Mechanical ventilation (among infants with home medical care)	 <p>1: Yes 19% 2: No 81%</p>	204

No.	Resources of participating hospitals	All hospitals	n
2352	Tracheostomy (among infants with home medical care)	 <p>1: Yes 20% 2: No 80%</p>	205
2353	Tube feeding (among infants with home medical care)	 <p>1: Yes 47% 2: No 53%</p>	211
2354	VP shunt (among infants with home medical care)	 <p>1: Yes 36% 2: No 64%</p>	204
2360	Rehabilitation (among infants with followup at 3 years of age)	 <p>1: Yes 22% 2: No 78%</p>	5287
2370	Behavioral disorder (among infants with followup at 3 years of age)	 <p>1: Yes 13% 2: No 87%</p>	5356
2372	Type of behavioral disorder (among infants with followup at 3 years of age)	 <p>1: ADHD or suspected ADHD 33% 2: ASD or suspected ASD 41% 3: Others 26%</p>	719