


















No.	Resources of participating hospitals	All hospitals	n
Q			
2210	Followup at 3 years of age (among infants with alive at discharge)	 <p>1:Yes 40% 2:No 60%</p>	68959
2212	Dead after discharge (among infants with alive at discharge)	 <p>1:Yes 1% 2:No 86% 3:not available 13%</p>	18730
2216	Reason for dropout (among infants with alive at discharge)	 <p>1:Followed at different hospital 61% 2:Admitted in rehabilitation center 1% 3:No contact 25% 4:Others 14% 5:Admitted in other rehabilitation center 0% 6:Hospitalization 0%</p>	7121
2220	Age at followup (mean) (among infants with followup at 3 years of age)	3.1	27138
	SD	0.2	
	95% confidence interval	3.1-3.1	
2222	Age corrected at followup (mean) (among infants with followup at 3 years of age)	2.4	11150
	SD	0.5	
	95% confidence interval	2.4-2.4	
2230	Body weight (mean) (among infants with followup at 3 years of age)	12.2	26779
	SD	1.8	
	95% confidence interval	12.1-12.2	







No.	Resources of participating hospitals	All hospitals	n
2240	Height (mean) (among infants with followup at 3 years of age)	89.4	26427
	SD	4.3	
	95% confidence interval	89.3-89.4	
2250	Head circumference (mean) (among infants with followup at 3 years of age)	48.3	23873
	SD	2.0	
	95% confidence interval	48.3-48.3	
2252	Chest circumference (mean) (among infants with followup at 3 years of age)	48.7	13807
	SD	2.7	
	95% confidence interval	48.7-48.8	
2254	Abdominal circumference (mean) (among infants with followup at 3 years of age)	46.2	6402
	SD	3.4	
	95% confidence interval	46.2-46.3	
2260	Oxygen (among infants with followup at 3 years of age)	<p>1: Yes 2% 2: No 98%</p>	22929
2261	Duration of home oxygen (mean) (among infants with oxygen)	17.1	444
	SD	9.4	
	95% confidence interval	16.2-17.9	







No.	Resources of participating hospitals	All hospitals	n
2270	Visual impairment (among infants with followup at 3 years of age)	 <p>1:Yes 6% 2:No 94%</p>	25740
2271	Severety of visual impairment (among infants with visual impairment)	 <p>1:Less than light perception 3% 2:Amblyopia or ny stagmus 28% 3:strabismus 32% 4:Others 37%</p>	631
2272	Eye glasses (among infants with visual impairment)	 <p>1:Yes 4% 2:No 96%</p>	10583
2280	Cerebral palsy (among infants with followup at 3 years of age)	 <p>1:Yes 8% 2:No 92%</p>	26667
2281	GMFCS grade (among infants with cerebral palsy)	 <p>1:I 25% 2:II 19% 3:III 14% 4:IV 17% 5:V 24%</p>	360
2282	Type of cerebral palsy (among infants with cerebral palsy)	 <p>1:Spastic 78% 2:Athetoid 4% 3:Mixed 5% 4:Flaccid 7% 5:others 6%</p>	609

No.	Resources of participating hospitals	All hospitals	n
2283	Cause of cerebral palsy (among infants with cerebral palsy)	 <p>1: PVL 53% 2: IVH 27% 3: Others 20%</p>	556
2285	DQ measurement (among infants with followup at 3 years of age)	 <p>1: Yes 91% 2: No 9%</p>	26593
2288	Reason not to measure DQ (among infants with DQ measurement)	 <p>1: Normal development by physician diagnosis 43% 2: Severely damaged by physician diagnosis 21% 3: Refusal from parents 4% 4: Impossible to perform due to severely damaged 8% 5: Failed to perform 11% 6: others 13%</p>	2325
2300	Method for DQ measurement (among infants with followup at 1.5 years of age)	 <p>1: Kyoto scale 88% 2: Others 12%</p>	24617
2301	DQ (K scale) (mean) (among infants with DQ measured by K scale)	83.7	21101
	SD	16.1	
	95% confidence interval	83.5-83.9	
2302	DQ corrected age (K scale) (mean) (among infants with DQ measured by K scale)	88.8	12322
	SD	17.7	
	95% confidence interval	88.5-89.1	

No.	Resources of participating hospitals	All hospitals	n
2303	DQ postural-motor (K scale) (mean) (among infants with DQ measured by K scale)	86.2	16396
	SD	21.2	
	95% confidence interval	85.9-86.5	
2304	DQ postural-motor corrected age (K scale) (mean) (among infants with DQ measured by K scale)	91.3	12229
	SD	22.9	
	95% confidence interval	90.9-91.7	
2305	DQ cognitive-adaptive (K scale) (mean) (among infants with DQ measured by K scale)	84.6	16537
	SD	16.6	
	95% confidence interval	84.3-84.8	
2306	DQ cognitive-adaptive corrected age (K scale) (mean) (among infants with DQ measured by K scale)	89.9	12313
	SD	18.2	
	95% confidence interval	89.6-90.3	
2307	DQ language-social (K scale) (mean) (among infants with DQ measured by K scale)	82.6	16514
	SD	18.5	
	95% confidence interval	82.3-82.9	
2308	DQ language-social corrected age (K scale) (mean) (among infants with DQ measured by K scale)	87.9	12270
	SD	20.0	
	95% confidence interval	87.5-88.2	

No.	Resources of participating hospitals	All hospitals	n
2311	Method for DQ other than K scale (among infants with DQ measured by other than K scale)	 <p>1: Bayley 7% 2: Enjogi 63% 3: Tsumori-Inage 18% 4: Others 12%</p>	2376
2312	DQ (other than K scale) (mean) (among infants with DQ measured by other than K scale)	89.3	2367
	SD	22.2	
	95% confidence interval	88.4-90.2	
2313	DQ corrected age (other than K scale) (mean) (among infants with DQ measured by other than K scale)	98.1	945
	SD	22.6	
	95% confidence interval	96.7-99.5	
2314	Evaluation (other than K scale) (among infants with DQ measured by other than K scale)	 <p>1: Normal 75% 2: Bordeline 11% 3: Delayed 13%</p>	824
2315	Evaluation by physician (among infants with DQ measured by other than K scale)	 <p>1: Normal 75% 2: Bordeline 9% 3: Delayed 16%</p>	558
2320	Hearing impairment (among infants with followup at 3 years of age)	 <p>1: Yes 2% 2: No 98%</p>	10825

No.	Resources of participating hospitals	All hospitals	n
2322	Hearing aid (among infants with hearing impairment)	 <p>1: Yes 46% 2: No 54%</p>	181
2323	Audiometry (among infants with hearing impairment)	 <p>1: Normal 26% 2: Moderate 53% 3: Severe 21%</p>	145
2330	Asthme (among infants with followup at 3 years of age)	 <p>1: Yes 8% 2: No 92%</p>	10833
2340	Epilepsy (among infants with followup at 3 years of age)	 <p>1: Yes 2% 2: No 98%</p>	11082
2350	Home medical care (among infants with followup at 3 years of age)	 <p>1: Yes 4% 2: No 96%</p>	9547
2351	Mechanical ventilation (among infants with home medical care)	 <p>1: Yes 16% 2: No 84%</p>	313

No.	Resources of participating hospitals	All hospitals	n
2352	Tracheostomy (among infants with home medical care)	 <p>1: Yes 22% 2: No 78%</p>	316
2353	Tube feeding (among infants with home medical care)	 <p>1: Yes 48% 2: No 52%</p>	330
2354	VP shunt (among infants with home medical care)	 <p>1: Yes 32% 2: No 68%</p>	315
2360	Rehabilitation (among infants with followup at 3 years of age)	 <p>1: Yes 23% 2: No 77%</p>	9501
2370	Behavioral disorder (among infants with followup at 3 years of age)	 <p>1: Yes 12% 2: No 88%</p>	9425
2372	Type of behavioral disorder (among infants with followup at 3 years of age)	 <p>1: ADHD or suspected ADHD 30% 2: ASD or suspected ASD 47% 3: Others 23%</p>	1155