



















No.	Resources of participating hospitals	All hospitals	n
Q			
2210	Followup at 3 years of age (among infants with alive at discharge)	 <p>1:Yes 39% 2:No 61%</p>	83568
2212	Dead after discharge (among infants with alive at discharge)	 <p>1:Yes 1% 2:No 99% 3:not available 0%</p>	16952
2216	Reason for dropout (among infants with alive at discharge)	 <p>1:Followed at different hospital 61% 2:Admitted in rehabilitation center 1% 3:No contact 25% 4:Others 14% 5:Admitted in other rehabilitation center 0% 6:Hospitalization 0%</p>	8294
2220	Age at followup (mean) (among infants with followup at 3 years of age)	3.1	31843
	SD	0.2	
	95% confidence interval	3.1-3.1	
2222	Age corrected at followup (mean) (among infants with followup at 3 years of age)	2.4	15855
	SD	0.5	
	95% confidence interval	2.4-2.4	
2230	Body weight (mean) (among infants with followup at 3 years of age)	12.2	31319
	SD	1.8	
	95% confidence interval	12.2-12.2	







No.	Resources of participating hospitals	All hospitals	n
2240	Height (mean) (among infants with followup at 3 years of age)	89.4	30891
	SD	4.3	
	95% confidence interval	89.3-89.4	
2250	Head circumference (mean) (among infants with followup at 3 years of age)	48.3	27795
	SD	2.0	
	95% confidence interval	48.3-48.3	
2252	Chest circumference (mean) (among infants with followup at 3 years of age)	48.7	16036
	SD	2.7	
	95% confidence interval	48.7-48.8	
2254	Abdominal circumference (mean) (among infants with followup at 3 years of age)	46.3	7734
	SD	3.4	
	95% confidence interval	46.2-46.4	
2260	Oxygen (among infants with followup at 3 years of age)	 <p>1: Yes 3% 2: No 97%</p>	27256
2261	Duration of home oxygen (mean) (among infants with oxygen)	17.2	638
	SD	9.2	
	95% confidence interval	16.5-17.9	







No.	Resources of participating hospitals	All hospitals	n
2270	Visual impairment (among infants with followup at 3 years of age)	 <p>1:Yes 6% 2:No 94%</p>	30049
2271	Severety of visual impairment (among infants with visual impairment)	 <p>1:Less than light perception 2% 2:Amblyopia or ny stagmus 29% 3:strabismus 32% 4:Others 37%</p>	904
2272	Eye glasses (among infants with visual impairment)	 <p>1:Yes 4% 2:No 96%</p>	15006
2280	Cerebral palsy (among infants with followup at 3 years of age)	 <p>1:Yes 7% 2:No 93%</p>	31077
2281	GMFCS grade (among infants with cerebral palsy)	 <p>1:I 24% 2: II 19% 3:III 14% 4:IV 17% 5: V 26%</p>	523
2282	Type of cerebral palsy (among infants with cerebral palsy)	 <p>1:Spastic 78% 2:Athetoid 3% 3:Mixed 6% 4:Flaccid 7% 5:others 6%</p>	802

No.	Resources of participating hospitals	All hospitals	n
2283	Cause of cerebral palsy (among infants with cerebral palsy)	 <p>1:PVL 54% 2:IVH 26% 3:Others 19%</p>	756
2285	DQ measurement (among infants with followup at 3 years of age)	 <p>1:Yes 91% 2:No 9%</p>	31288
2288	Reason not to measure DQ (among infants with DQ measurement)	 <p>1:Normal development by physician diagnosis 43% 2:Severely damaged by physician diagnosis 22% 3:Refusal from patients 4% 4:Impossible to perform due to severely damaged 7% 5:Failed to perform 10% 6:others 14%</p>	2683
2300	Method for DQ measurement (among infants with followup at 1.5 years of age)	 <p>1:Kyoto scale 88% 2:Others 12%</p>	28887
2301	DQ (K scale) (mean) (among infants with DQ measured by K scale)	83.6	24693
	SD	16.3	
	95% confidence interval	83.4-83.8	
2302	DQ corrected age (K scale) (mean) (among infants with DQ measured by K scale)	88.7	15384
	SD	17.6	
	95% confidence interval	88.4-88.9	

No.	Resources of participating hospitals	All hospitals	n
2303	DQ postural-motor (K scale) (mean) (among infants with DQ measured by K scale)	86.3	20080
	SD	21.2	
	95% confidence interval	86.0-86.6	
2304	DQ postural-motor corrected age (K scale) (mean) (among infants with DQ measured by K scale)	91.5	15283
	SD	22.6	
	95% confidence interval	91.2-91.9	
2305	DQ cognitive-adaptive (K scale) (mean) (among infants with DQ measured by K scale)	84.4	20222
	SD	16.8	
	95% confidence interval	84.2-84.6	
2306	DQ cognitive-adaptive corrected age (K scale) (mean) (among infants with DQ measured by K scale)	89.8	15378
	SD	18.1	
	95% confidence interval	89.5-90.0	
2307	DQ language-social (K scale) (mean) (among infants with DQ measured by K scale)	82.5	20199
	SD	18.7	
	95% confidence interval	82.2-82.7	
2308	DQ language-social corrected age (K scale) (mean) (among infants with DQ measured by K scale)	87.8	15329
	SD	20.0	
	95% confidence interval	87.4-88.1	

No.	Resources of participating hospitals	All hospitals	n
2311	Method for DQ other than K scale (among infants with DQ measured by other than K scale)	 <p>1: Bayley 9% 2: Enjogi 63% 3: Tsumori-Inage 16% 4: Others 12%</p>	2700
2312	DQ (other than K scale) (mean) (among infants with DQ measured by other than K scale)	89.2	2543
	SD	22.3	
	95% confidence interval	88.3-90.0	
2313	DQ corrected age (other than K scale) (mean) (among infants with DQ measured by other than K scale)	96.9	1093
	SD	23.5	
	95% confidence interval	95.5-98.3	
2314	Evaluation (other than K scale) (among infants with DQ measured by other than K scale)	 <p>1: Normal 71% 2: Bordeline 12% 3: Delayed 17%</p>	1120
2315	Evaluation by physician (among infants with DQ measured by other than K scale)	 <p>1: Normal 75% 2: Bordeline 9% 3: Delayed 16%</p>	553
2320	Hearing impairment (among infants with followup at 3 years of age)	 <p>1: Yes 2% 2: No 98%</p>	15229

No.	Resources of participating hospitals	All hospitals	n
2322	Hearing aid (among infants with hearing impairment)	 <p>1:Yes 46% 2:No 54%</p>	248
2323	Audiometry (among infants with hearing impairment)	 <p>1:Normal 22% 2:Moderate 58% 3:Severe 20%</p>	187
2330	Asthme (among infants with followup at 3 years of age)	 <p>1:Yes 9% 2:No 91%</p>	15201
2340	Epilepsy (among infants with followup at 3 years of age)	 <p>1:Yes 2% 2:No 98%</p>	15535
2350	Home medical care (among infants with followup at 3 years of age)	 <p>1:Yes 4% 2:No 96%</p>	13896
2351	Mechanical ventilation (among infants with home medical care)	 <p>1:Yes 18% 2:No 82%</p>	438

No.	Resources of participating hospitals	All hospitals	n
2352	Tracheostomy (among infants with home medical care)	 <p>1:Yes 25% 2:No 75%</p>	441
2353	Tube feeding (among infants with home medical care)	 <p>1:Yes 51% 2:No 49%</p>	461
2354	VP shunt (among infants with home medical care)	 <p>1:Yes 30% 2:No 70%</p>	448
2360	Rehabilitation (among infants with followup at 3 years of age)	 <p>1:Yes 23% 2:No 77%</p>	13957
2370	Behavioral disorder (among infants with followup at 3 years of age)	 <p>1:Yes 12% 2:No 88%</p>	13769
2372	Type of behavioral disorder (among infants with followup at 3 years of age)	 <p>1:ADHD or suspected ADHD 30% 2:ASD or suspected ASD 47% 3:Others 23%</p>	1681