


















No.	Resources of participating hospitals	All hospitals	n
Q			
2210	Followup at 3 years of age (among infants with alive at discharge)	 1:Yes 39% 2:No 61%	86535
2212	Dead after discharge (among infants with alive at discharge)	 1:Yes 1% 2:No 99% 3:not available 0%	17001
2216	Reason for dropout (among infants with alive at discharge)	 1:Followed at different hospital 60% 2:Admitted in rehabilitation center 1% 3:No contact 25% 4:Others 14% 5:Admitted in other rehabilitation center 0% 6:Hospitalization 0%	8612
2220	Age at followup (mean) (among infants with followup at 3 years of age)	3.1	33303
	SD	0.2	
	95% confidence interval	3.1-3.1	
2222	Age corrected at followup (mean) (among infants with followup at 3 years of age)	2.4	17382
	SD	0.5	
	95% confidence interval	2.4-2.4	
2230	Body weight (mean) (among infants with followup at 3 years of age)	12.2	32712
	SD	1.8	
	95% confidence interval	12.2-12.2	







No.	Resources of participating hospitals	All hospitals	n
2240	Height (mean) (among infants with followup at 3 years of age)	89.4	32262
	SD	4.3	
	95% confidence interval	89.4-89.4	
2250	Head circumference (mean) (among infants with followup at 3 years of age)	48.3	28950
	SD	2.0	
	95% confidence interval	48.3-48.3	
2252	Chest circumference (mean) (among infants with followup at 3 years of age)	48.7	16702
	SD	2.7	
	95% confidence interval	48.7-48.8	
2254	Abdominal circumference (mean) (among infants with followup at 3 years of age)	46.3	8112
	SD	3.4	
	95% confidence interval	46.2-46.4	
2260	Oxygen (among infants with followup at 3 years of age)	1:Yes 3% 2:No 97%	28575
2261	Duration of home oxygen (mean) (among infants with oxygen)	17.2	700
	SD	9.1	
	95% confidence interval	16.5-17.9	

No.	Resources of participating hospitals	All hospitals	n
2270	Visual impairment (among infants with followup at 3 years of age)	 1:Yes 6% 2:No 94%	31371
2271	Severity of visual impairment (among infants with visual impairment)	 1:Less than light perception 2% 2:Amblyopia or nystagmus 29% 3:strabismus 32% 4:Others 37%	1012
2272	Eye glasses (among infants with visual impairment)	 1:Yes 4% 2:No 96%	16351
2280	Cerebral palsy (among infants with followup at 3 years of age)	 1:Yes 7% 2:No 93%	32413
2281	GMFCS grade (among infants with cerebral palsy)	 1:I 25% 2:II 19% 3:III 14% 4:IV 17% 5:V 25%	557
2282	Type of cerebral palsy (among infants with cerebral palsy)	 1:Spastic 78% 2:Athetoid 3% 3:Mixed 6% 4:Flaccid 7% 5:others 6%	852

No.	Resources of participating hospitals	All hospitals	n
2283	Cause of cerebral palsy (among infants with cerebral palsy)	 1:PVL 54% 2:IVH 26% 3:Others 20%	810
2285	DQ measurement (among infants with followup at 3 years of age)	 1:Yes 91% 2:No 9%	32751
2288	Reason not to measure DQ (among infants with DQ measurement)	 1:Normal development by physician diagnosis 43% 2:Severely damaged by physician diagnosis 22% 3:Refusal from parents 4% 4:Impossible to perform due to severely damaged 7% 5:Failed to perform 10% 6:others 13%	2760
2300	Method for DQ measurement (among infants with followup at 1.5 years of age)	 1:Kyoto scale 89% 2:Others 11%	30245
2301	DQ (K scale) (mean) (among infants with DQ measured by K scale)	83.5	25859
	SD	16.3	
	95% confidence interval	83.3-83.7	
2302	DQ corrected age (K scale) (mean) (among infants with DQ measured by K scale)	88.7	16375
	SD	17.6	
	95% confidence interval	88.4-88.9	

No.	Resources of participating hospitals	All hospitals	n
2303	DQ postural-motor (K scale) (mean) (among infants with DQ measured by K scale)	86.2	21223
	SD	21.2	
	95% confidence interval	85.9-86.5	
2304	DQ postural-motor corrected age (K scale) (mean) (among infants with DQ measured by K scale)	91.5	16273
	SD	22.6	
	95% confidence interval	91.2-91.9	
2305	DQ cognitive-adaptive (K scale) (mean) (among infants with DQ measured by K scale)	84.4	21363
	SD	16.9	
	95% confidence interval	84.2-84.6	
2306	DQ cognitive-adaptive corrected age (K scale) (mean) (among infants with DQ measured by K scale)	89.8	16367
	SD	18.2	
	95% confidence interval	89.5-90.0	
2307	DQ language-social (K scale) (mean) (among infants with DQ measured by K scale)	82.4	21342
	SD	18.8	
	95% confidence interval	82.2-82.7	
2308	DQ language-social corrected age (K scale) (mean) (among infants with DQ measured by K scale)	87.8	16321
	SD	20.0	
	95% confidence interval	87.5-88.1	

No.	Resources of participating hospitals	All hospitals	n
2311	Method for DQ other than K scale (among infants with DQ measured by other than K scale)	 1:Bayley 9% 2:Enjogi 63% 3:Tsumori-Inage 16% 4:Others 12%	2792
2312	DQ (other than K scale) (mean) (among infants with DQ measured by other than K scale)	89.2	2585
	SD	22.3	
	95% confidence interval	88.3-90.1	
2313	DQ corrected age (other than K scale) (mean) (among infants with DQ measured by other than K scale)	97.1	1125
	SD	23.2	
	95% confidence interval	95.7-98.4	
2314	Evaluation (other than K scale) (among infants with DQ measured by other than K scale)	 1:Normal 70% 2:Borderline 13% 3:Delayed 17%	1204
2315	Evaluation by physician (among infants with DQ measured by other than K scale)	 1:Normal 75% 2:Borderline 9% 3:Delayed 16%	551
2320	Hearing impairment (among infants with followup at 3 years of age)	 1:Yes 2% 2:No 98%	16613

No.	Resources of participating hospitals	All hospitals	n
2322	Hearing aid (among infants with hearing impairment)	 1:Yes 45% 2:No 55%	261
2323	Audiometry (among infants with hearing impairment)	 1:Normal 22% 2:Moderate 58% 3:Severe 20%	199
2330	Asthme (among infants with followup at 3 years of age)	 1:Yes 9% 2:No 91%	16593
2340	Epilepsy (among infants with followup at 3 years of age)	 1:Yes 2% 2:No 98%	16934
2350	Home medical care (among infants with followup at 3 years of age)	 1:Yes 4% 2:No 96%	15268
2351	Mechanical ventilation (among infants with home medical care)	 1:Yes 19% 2:No 81%	467

No.	Resources of participating hospitals	All hospitals	n
2352	Tracheostomy (among infants with home medical care)	1: Yes 25% 2: No 75%	471
2353	Tube feeding (among infants with home medical care)	1: Yes 53% 2: No 47%	498
2354	VP shunt (among infants with home medical care)	1: Yes 29% 2: No 71%	479
2360	Rehabilitation (among infants with followup at 3 years of age)	1: Yes 23% 2: No 77%	15351
2370	Behavioral disorder (among infants with followup at 3 years of age)	1: Yes 12% 2: No 88%	15122
2372	Type of behavioral disorder (among infants with followup at 3 years of age)	1: ADHD or suspected ADHD 29% 2: ASD or suspected ASD 47% 3: Others 24%	1819