

1. Tachikawa T, Masunaga K, Kiyota M, Kushima R, Nishina S, Isayama T, Noda E, Okazaki K, Funaki T, Nakao A, Matsumoto T, Arai H, Saito Y, Miyazawa T, Negishi T, Shoji H, Toyama T, Nishimura R, Ueno R, Sato F, Morikawa Y; Neonatal Research Network of Japan. Ten-year changes in risk factors for requiring treatment of retinopathy of prematurity in extremely low birth weight infants: a multicenter study in Tokyo. *Jpn J Ophthalmol.* 2026 Jan 14. doi: 10.1007/s10384-025-01322-z.
2. Katsuki S, Ushida T, Nakatochi M, Kobayashi Y, Tano S, Matsuo S, Imai K, Sato Y, Hayakawa M, Kajiyama H, Kotani T; Neonatal Research Network of Japan. Hypertensive disorders of pregnancy and retinopathy of prematurity in extremely preterm infants: A causal mediation analysis. *Pregnancy Hypertens.* 2026 Jan 7;43:101410. doi: 10.1016/j.preghy.2025.101410.
3. Ishida S, Nakanishi H, Sekiya R, Kawada K, Kosaka Y, Yamaguchi A, Ooka M; Neonatal Research Network, Japan. Impact of chorionicity and histological chorioamnionitis on neurodevelopment and mortality in extremely preterm twins. *Early Hum Dev.* 2025;210:106370. doi: 10.1016/j.earlhumdev.2025.106370.
4. Isayama T, Norman M, Kusuda S, Reichman B, Lehtonen L, Lui K, Adams M, Vento Torres M, Filippi L, Battin M, Guinsburg R, Modi N, Håkansson S, Klinger G, de Almeida MF, Helenius K, Bassler D, Su YC, Shah PS; International Network for Evaluation of Outcomes (iNeo) Investigators. Outcomes of Preterm Infants Born at 22 to 23 Weeks' Gestation in 11 International Neonatal Networks. *JAMA Pediatr.* 2025;179(11):1183-1193. doi: 10.1001/jamapediatrics.2025.2958.
5. Hollens G, Schindler T, Battin M, Klinger G, Adams M, Vento M, Santacroce A, Håkansson S, Isayama T, Norman M, Kusuda S, Lehtonen L, Helenius K, Modi N, Shah PS, Lui K; iNeo Neonatal Network. International Variation and Trends of Intraventricular Hemorrhage in Very Preterm Infants. *Neonatology.* 2025;122(5):570-585. doi: 10.1159/000546714.
6. Klinger G, Helenius K, Vento M, Kusuda S, Norman M, Procianoy RS, Goswami N, Biran V, Bassler D, Reichman B, Skubisz A, Battin M, Lehtonen L, Lui K, Mori A, Beltempo M, Adams M, San Feliciano L, Isayama T, Shah PS; iNeo Investigators. Transfusion Practices in 12 Neonatal Networks: Are We Closer to Adopting a Restrictive Transfusion Approach? *Neonatology.* 2025;122(5):561-569. doi: 10.1159/000546612.
7. Namba F, Nakanishi H. Shifting Paradigms in Bronchopulmonary Dysplasia: From Treatment to Etiology/Pathophysiology-Based Classification. *Biomedicines.* 2025;13(4):985. doi: 10.3390/biomedicines13040985.
8. Kusuda S, Bennett MV, Gould JB; Neonatal Research Network of Japan. Comparative analysis of necrotizing enterocolitis in preterm infants born in Japan and born to mothers of Japanese ethnicity in California. *Sci Rep.* 2025;15(1):9943. doi: 10.1038/s41598-025-92393-y.

9. Ishida S, Nakanishi H, Sekiya R, Kawada K, Kosaka Y, Yamaguchi A, Ooka M; Neonatal Research Network, Japan. Impact of late-onset circulatory collapse and bronchopulmonary dysplasia on short-term outcomes in preterm infants: A Japanese retrospective cohort study. *Pediatr Neonatol.* 2025;66(5):462-466. doi: 10.1016/j.pedneo.2024.08.006.
10. Kanzawa T, Kinoshita F, Namba F, Tanaka T, Oshiro M, Sugiura T, Kato Y, Miyata M, Yamada Y, Iwata O, Hayakawa M, Sato Y; BPD Prediction Model Collaborative Clinical Research Team; Neonatal Research Network of Japan. Development and validation of a prediction model for bronchopulmonary dysplasia using respiratory severity score. *Pediatr Res.* 2025;98(2):577-584. doi: 10.1038/s41390-025-03862-z.
11. Nakanishi H, Ito M, Kato S, Saito M, Miyahara N, Arai H, Ota E, Namba F; Neonatal Research Network, Japan. Validation of a New Classification for Severe Bronchopulmonary Dysplasia in Extremely Preterm Infants: Insights from a Large Japanese Cohort. *Neonatology.* 2025;122(3):339-349. doi: 10.1159/000543810.
12. Fenton TR, Alshaikh B, Kusuda S, Helenius K, Modi N, Norman M, Lui K, Lehtonen L, Battin M, Klinger G, Vento M, Lastrucci V, Gagliardi L, Adams M, Marba STM, Isayama T, Hakansson S, Bassler D, Shah PS; International Network for Evaluation of Outcomes (iNeo) of Neonates Investigators. Birth weight and head circumference for 22-29 weeks gestation neonates from an international cohort. *Arch Dis Child Fetal Neonatal Ed.* 2025;110(4):401-408. doi: 10.1136/archdischild-2024-327845.
13. Ushida T, Nosaka R, Nakatochi M, Kobayashi Y, Tano S, Fuma K, Matsuo S, Imai K, Sato Y, Hayakawa M, Kajiyama H, Kotani T; Neonatal Research Network of Japan. Effect of chorioamnionitis on postnatal growth in very preterm infants: a population-based study in Japan. *Arch Gynecol Obstet.* 2025;311(5):1321-1330. doi: 10.1007/s00404-024-07757-y.
14. Klinger G, Reichman B, Norman M, Kusuda S, Battin M, Helenius K, Isayama T, Lui K, Adams M, Vento M, Hakansson S, Beltempo M, Poggi C, San Feliciano L, Lehtonen L, Bassler D, Yang J, Shah PS; International Network for Evaluating Outcomes of Neonates (iNeo). Late-Onset Sepsis among Extremely Preterm Infants of 24-28 Weeks Gestation: An International Comparison in 10 High-Income Countries. *Neonatology.* 2024;121(6):761-771. doi: 10.1159/000539245.
15. Sugie M, Nawa N, Noguchi Y, Taki A, Kashimada A, Honda I, Koyama A, Okazaki K, Kondo M, Miyahara H, Ito K, Yamauchi T, Kondo T, Honda-Ozaki F, Kusuda S, Morioka C, Fujiwara T, Morio T, Kashimada K. Stage III Chorioamnionitis is Associated with Reduced Risk of Severe Retinopathy of Prematurity. *J Pediatr.* 2024;272:114085. doi: 10.1016/j.jpeds.2024.114085.
16. Yoneda K, Seki T, Kawazoe Y, Ohe K, Takahashi N; Neonatal Research Network of Japan. Immediate postnatal prediction of death or

bronchopulmonary dysplasia among very preterm and very low birth weight infants based on gradient boosting decision trees algorithm: A nationwide database study in Japan. *PLoS One*. 2024;19(3):e0300817. doi: 10.1371/journal.pone.0300817.

17. Fukui K, Ito Y, Kokubo M, Nakanishi H, Hirano S, Kusuda S, Ito S, Isayama T; Neonatal Research Network of Japan. Erythropoietin and retinopathy of prematurity: a retrospective cohort study in Japan, 2008-2018. *J Perinatol*. 2024;44(6):886-891. doi: 10.1038/s41372-024-01929-z.
18. Tamai K, Matsumoto N, Yorifuji T, Takeuchi A, Nakamura M, Nakamura K, Kageyama M; Neonatal Research Network of Japan. Delivery room intubation and neurodevelopment among extremely preterm infants. *Pediatr Res*. 2024;95(7):1897-1902. doi: 10.1038/s41390-023-02993-5.
19. Miyake H, Nakano R, Yamamoto S, Isayama T, Sasaki H; Neonatal Research Network of Japan. Mortality and neurodevelopmental outcomes in very low birth weight infants with esophageal atresia. *Pediatr Surg Int*. 2023;39(1):294. doi: 10.1007/s00383-023-05579-8.
20. Tamai K, Matsumoto N, Yorifuji T, Takeuchi A, Nakamura M, Nakamura K, Kageyama M; Neonatal Research Network of Japan. Delivery room intubation and severe intraventricular hemorrhage in extremely preterm infants without low Apgar scores: A Japanese retrospective cohort study. *Sci Rep*. 2023;13(1):14990. doi: 10.1038/s41598-023-41010-x.
21. Inoue T, Nishikubo T, Hirano S, Kamamoto T, Takahashi Y, Kusuda S; Neonatal Research Network of Japan. Risk factor analyses for intraventricular hemorrhage in preterm infants: A retrospective cohort study. *Pediatr Int*. 2023;65(1):e15599. doi: 10.1111/ped.15599.
22. Parikh S, Reichman B, Kusuda S, Adams M, Lehtonen L, Vento M, Norman M, San Feliciano L, Isayama T, Hakansson S, Helenius K, Bassler D, Yang J, Shah PS, Gellineo L; International Network for Evaluation of Outcomes (iNeo) of neonates investigators. Trends, Characteristic, and Outcomes of Preterm Infants Who Received Postnatal Corticosteroid: A Cohort Study from 7 High-Income Countries. *Neonatology*. 2023 Apr 25:1-10. Ishida S, Nakanishi H, Kosaka Y, Yamaguchi A, Ooka M; Neonatal Research Network, Japan. Evaluation of newborns with Down syndrome with weight less than 1500g in the neonatal intensive care unit: A Japanese multicentre study. *J Paediatr Child Health*. 2023;59:912-918.
23. Ishida S, Nakanishi H, Sekiya R, Kawada K, Kosaka Y, Yamaguchi A, Ooka M, On behalf of the Neonatal Research Network of Japan. Evaluation of Postnatal Complications in Clinical and Histological Chorioamnionitis in Extremely Preterm Infants: A Japanese Cohort Study. *Am J Perinatol* 2023 DOI: 10.1055/a-2113-4332.
24. Ishida S, Nakanishi H, Kosaka Y, Yamaguchi A, Ooka M; Neonatal

- Research Network, Japan. Evaluation of newborns with Down syndrome with weight less than 1500 g in the neonatal intensive care unit: A Japanese multicentre study. *J Paediatr Child Health*. 2023;59(7):912-918. doi: 10.1111/jpc.16418.
25. Parikh S, Reichman B, Kusuda S, Adams M, Lehtonen L, Vento M, Norman M, San Feliciano L, Isayama T, Hakansson S, Helenius K, Bassler D, Yang J, Shah PS, Gellineo L; International Network for Evaluation of Outcomes (iNeo) of neonates investigators. Trends, Characteristic, and Outcomes of Preterm Infants Who Received Postnatal Corticosteroid: A Cohort Study from 7 High-Income Countries. *Neonatology*. 2023 Apr 25:1-10. doi: 10.1159/000530128
 26. Kusuda S, Nakanishi H, Suzuki H; Neonatal Research Network of Japan. Validation of eligibility criteria for the Japan Obstetric Compensation System for Cerebral Palsy in preterm infants: A case-control cohort study. *J Obstet Gynaecol Res*. 2023;49:220-231.
 27. Nakanishi H, Isayama T, Kokubo M, Hirano S, Kusuda S; Neonatal Research Network, Japan. Inhaled Nitric Oxide Therapy in the Post-Acute Phase in Extremely Preterm Infants: A Japanese Cohort Study. *J Pediatr*. 2023;252:61-67.e5.
 28. Suenaga H, Nakanishi H, Uchiyama A, Kusuda S; Neonatal Research Network of Japan. Small for Gestational Age Affects Outcomes on Singletons and Inborn Births in Extremely Preterm Infants: A Japanese Cohort Study. *Am J Perinatol*. 2022: 10.1055/a-1933-4627.
 29. Kawasaki H, Yamada T, Takahashi Y, Nakayama T, Wada T, Kosugi S, Neonatal Research Network of Japan. Mortality and morbidity of infants with trisomy 21, weighing 1500 grams or less, in Japan. *J Hum Genet* . 2022 Jul 5. doi: 10.1038/s10038-022-01061-w.
 30. Kusuda S, Masumoto K, Nakanishi H. Care of periviable preterm infants—Unique challenges from Japan. *Neonatal-Perinatal Medicine* (12 Ed). 2023
 31. Kusuda S, Nakanishi H, Oei JL. Practical procedures for the delivery room resuscitation of micropreemies. *Semin Fetal Neonatal Med*. 2022;27:101349.
 32. Yanagisawa T, Nakamura T, Kokubo M. Prognosis of 22- and 23-Gestational-Week-Old Infants at Our Facility: A Retrospective Cohort Study. *Am J Perinatol*. 2022. doi: 10.1055/a-1779-4032.
 33. Isayama T, Kusuda S, Adams M, Berti E, Battin M, Helenius K, Hakansson S, Vento M, Norman M, Reichman B, Noguchi A, Lee SK, Bassler D, Lui K, Lehtonen L, Yang J, Shah PS; International Network for Evaluating Outcomes of Neonates (iNeo) Investigators. International Variation in the Management of Patent Ductus Arteriosus and Its Association with Infant Outcomes: A Survey and Linked Cohort Study. *J Pediatr*. 2022. doi: 10.1016/j.jpeds.2021.12.071.
 34. Ushida T, Kotani T, Baba J, Imai K, Moriyama Y, Nakano-Kobayashi T, Iitani Y, Nakamura N, Hayakawa M, Kajiyama H; Neonatal Research

- Network of Japan. Antenatal prediction models for outcomes of extremely and very preterm infants based on machine learning. *Arch Gynecol Obstet*. 2022. doi: 10.1007/s00404-022-06865-x.
35. Ushida T, Nakatochi M, Kobayashi Y, Nakamura N, Fuma K, Iitani Y, Imai K, Sato Y, Hayakawa M, Kajiyama H, Kotani T; Neonatal Research Network of Japan. Antenatal corticosteroids and outcomes of small for gestational age infants born at 24-31 gestational weeks: a population-based propensity score matching analysis. *Arch Gynecol Obstet*. 2022. doi: 10.1007/s00404-022-06834-4.
 36. Ushida T, Nakamura N, Nakatochi M, Kobayashi Y, Sato Y, Iitani Y, Imai K, Nakano-Kobayashi T, Hayakawa M, Kajiyama H, Kotani T; Neonatal Research Network of Japan. Impact of hypertensive disorders of pregnancy on respiratory outcomes in extremely and very preterm infants: A population-based study in Japan. *Pregnancy Hypertens*. 2022;29:54-60.
 37. Ushida T, Kotani T, Nakatochi M, Kobayashi Y, Nakamura N, Imai K, Iitani Y, Nakano-Kobayashi T, Hayakawa M, Kajiyama H; Neonatal Research Network of Japan. Intrauterine exposure to hypertensive disorders of pregnancy and postnatal growth in extremely and very preterm infants. *Pregnancy Hypertens*. 2022;28:174-179.
 38. Suga S, Kyono Y, Kido T, Nakasone R, Abe S, Ashina M, Nozu K, Fujioka K. Long use of continuous positive airway pressure protects against the development of treatment-requiring retinopathy of prematurity. *Sci Rep*. 2022;12:7799.
 39. Yanagisawa T, Nakamura T, Kokubo M. Prognosis of 22- and 23-Gestational-Week-Old Infants at Our Facility: A Retrospective Cohort Study. *Am J Perinatol*. 2022. doi: 10.1055/a-1779-4032.
 40. Tomo CK, Balogun OO, Davidson J, Guinsburg R, Almeida MFB, Lopes JMA, Barros MCM, Takehara K, Mikami M, Isayama T, Hoshino A, Mori R, Mizuguchi M. Comparison of mortality and survival without major morbidities of very preterm infants with very low birth weight from Japan and Brazil. *Rev Paul Pediatr*. 2022;41:e2021389
 41. Kusuda S, Hirano S, Nakamura T. Creating experiences from active treatment towards extremely preterm infants born at less than 25 weeks in Japan. *Semin Perinatol*. 2022;46:151537.
 42. Kono Y; Neonatal Research Network of Japan. Neurodevelopmental outcomes of very low birth weight infants in the Neonatal Research Network of Japan: importance of neonatal intensive care unit graduate follow-up. *Clin Exp Pediatr*. 2021;64:313-321.
 43. Shah PS, Norman M, Rusconi F, Kusuda S, Reichman B, Battin M, Bassler D, Modi N, Hakansson S, Yang J, Lee SK, Helenius K, Vento M, Lehtonen L, Adams M, Isayama T, Lui K, Gagliardi L; International Network for Evaluating Outcomes of Neonates (iNeo) Investigators. Five-minute Apgar score and outcomes in neonates of 24-28 weeks' gestation. *Arch Dis Child Fetal Neonatal Ed*. 2021. doi:

10.1136/archdischild-2021-322230.

44. Kusuda S, Fujimura M. Neonatal network database operated by the Neonatal Research Network of Japan. *Pediatric Medicine* 2021. <https://dx.doi.org/10.21037/pm-21-71>
45. Seaton SE, Draper ES, Adams M, Kusuda S, Håkansson S, Helenius K, Reichman B, Lehtonen L, Bassler D, Lee SK, Vento M, Darlow BA, Rusconi F, Beltempo M, Isayama T, Lui K, Norman M, Yang J, Shah PS, Modi N; UK Neonatal Collaborative; International Network for Evaluating Outcomes of Neonates (iNeo) Investigators; ANZNN (Australian and New Zealand Neonatal Network); CNN (Canadian Neonatal Network); NRNJ (Neonatal Research Network Japan); SEN1500 (Spanish Neonatal Network); SwissNeoNet (Swiss Neonatal Network). Variations in Neonatal Length of Stay of Babies Born Extremely Preterm: An International Comparison Between iNeo Networks. *J Pediatr.* 2021;233:26-32.e6.
46. Nakamura N, Ushida T, Nakatochi M, Kobayashi Y, Moriyama Y, Imai K, Nakano-Kobayashi T, Hayakawa M, Kajiyama H, Kikkawa F, Kotani T, Neonatal Research Network of Japan. Mortality and neurological outcomes in extremely and very preterm infants born to mothers with hypertensive disorders of pregnancy. *Sci Rep* 2021;11:1729.
47. Kono Y. Neurodevelopmental outcomes of very low birth weight infants in the Neonatal Research Network of Japan: Importance of NICU graduate follow-up. *Clin Exp Pediatr.* 2021;64:313-321.
48. Nakashima T, Inoue H, Sakemi Y, Ochiai M, Yamashita H, Ohga S; Neonatal Research Network of Japan. Trends in Bronchopulmonary Dysplasia Among Extremely Preterm Infants in Japan, 2003-2016. *J Pediatr.* 2021;230:119-125.e7.
49. Sotiropoulos JX, Kapadia V, Vento M, Rabi Y, Saugstad OD, Kumar RK, Schmölzer GM, Zhang H, Yuan Y, Lim G, Kusuda S, Arimitsu T, Nguyen TT, Kitsommart R, Yeo KT, Oei JL. Oxygen for the delivery room respiratory support of moderate-to-late preterm infants. An international survey of clinical practice from 21 countries. *Acta Paediatr.* 2021;110:3261-3268.
50. Ushida T, Moriyama Y, Nakatochi M, Kobayashi Y, Imai K, Nakano-Kobayashi T, Nakamura N, Hayakawa M, Kajiyama H, Kotani T; Neonatal Research Network of Japan. Antenatal prediction models for short- and medium-term outcomes in preterm infants. *Acta Obstet Gynecol Scand.* 2021 Mar 3. doi: 10.1111/aogs.14136.
51. Kono Y, Rysavy MA, Nakanishi H, Kusuda S. Outcomes of infants born at periviable gestations in Japan. *Acta Paediatr.* 2021. doi: 10.1111/apa.15796.
52. Kono Y, Yonemoto N, Nakanishi H, Hosono S, Hirano S, Kusuda S, Fujimura M; Neonatal Research Network Japan. A Retrospective Cohort Study on Mortality and Neurodevelopmental Outcomes of Preterm Very Low Birth Weight Infants Born to Mothers with Hypertensive Disorders

- of Pregnancy. *Am J Perinatol*. 2021 Feb 3. doi: 10.1055/s-0041-1722874.
53. Lui K, Vento M, Modi N, Kusuda S, Lehtonen L, Håkansson S, Rusconi F, Bassler D, Reichman B, Yang J, Shah PS; International Network for Evaluating Outcomes of Neonates (iNeo) Investigators and Collaborators. Inter-center variability in neonatal outcomes of preterm infants: A longitudinal evaluation of 298 neonatal units in 11 countries. *Semin Fetal Neonatal Med*. 2021;26:101196.
 54. Murano Y, Shoji H, Ikeda N, Okawa N, Hayashi K, Kantake M, Morisaki N, Shimizu T, Gilmour S. Analysis of Factors Associated With Body Mass Index at Ages 18 and 36 Months Among Infants Born Extremely Preterm. *JAMA Netw Open*. 2021;4:e2128555.
 55. Rysavy MA, Mehler K, Oberthür A, Ågren J, Kusuda S, McNamara PJ, Giesinger RE, Kribs A, Normann E, Carlson SJ, Klein JM, Backes CH, Bell EF. An Immature Science: Intensive Care for Infants Born at ≤ 23 Weeks of Gestation. *J Pediatr*. 2021;233:16-25.e1.
 56. Ozawa J, Tanaka K, Kabe K, Namba F; Neonatal Research Network of Japan. Impact of being large-for-gestational-age on neonatal mortality and morbidities in extremely premature infants. *Pediatr Res*. 2021;1-7. doi: 10.1038/s41390-021-01375-z.
 57. Nakamura N, Ushida T, Nakatochi M, Kobayashi Y, Moriyama Y, Imai K, Nakano-Kobayashi T, Hayakawa M, Kajiyama H, Kikkawa F, Kotani T; Neonatal Research Network of Japan. Mortality and neurological outcomes in extremely and very preterm infants born to mothers with hypertensive disorders of pregnancy. *Sci Rep*. 2021;11:1729.
 58. Kusuda S, Bennett M, Gould J; Neonatal Research Network of Japan and the California Perinatal Quality Care Collaborative. Outcomes of Infants with Very Low Birth Weight Associated with Birthplace Difference: A Retrospective Cohort Study of Births in Japan and California. *J Pediatr*. 2021;229:182-190.e6. Kawasaki H, Yamada T, Takahashi Y, Nakayama T, Wada T, Kosugi S, Neonatal Research Network of Japan. Epidemiology of Birth Defects in Very Low Birth Weight Infants in Japan. *J Pediatr*. 2020;S0022-3476(20)30855-6.
 59. Helenius K, Morisaki N, Kusuda S, Shah PS, Norman M, Lehtonen L, Reichman B, Darlow BA, Noguchi A, Adams M, Bassler D, Håkansson S, Isayama T, Berti E, Lee SK, Vento M, Lui K; International Network for Evaluation of Outcomes of neonates (iNeo). Survey shows marked variations in approaches to redirection of care for critically ill very preterm infants in 11 countries. *Acta Paediatr*. 2020;109:1338-1345.
 60. Miyake F, Ito M, Minami H, Tamura M, Namba F. Management of bronchopulmonary dysplasia in Japan: A 10-year nationwide survey. *Pediatr Neonatol*. 2020;61:272-278.
 61. Norman M, Håkansson S, Kusuda S, Vento M, Lehtonen L, Reichman B, Darlow BA, Adams M, Bassler D, Isayama T, Rusconi F, Lee S, Lui K, Yang J, Shah PS; International Network for Evaluation of Outcomes in Neonates (iNeo) Investigators. Neonatal Outcomes in Very Preterm

- Infants With Severe Congenital Heart Defects: An International Cohort Study. *J Am Heart Assoc.* 2020;9:e015369.
62. Isayama T, Kusuda S, Reichman B, Lee SK, Lehtonen L, Norman M, Adams M, Bassler D, Helenius K, Hakansson S, Yang J, Jain A, Shah PS; International Network for Evaluating Outcomes of Neonates (iNeo) Investigators. Neonatal Intensive Care Unit-Level Patent Ductus Arteriosus Treatment Rates and Outcomes in Infants Born Extremely Preterm. *J Pediatr.* 2020;220:34-39.e5.
 63. Gagliardi L, Rusconi F, Reichman B, Adams M, Modi N, Lehtonen L, Kusuda S, Vento M, Darlow BA, Bassler D, Isayama T, Norman M, Håkansson S, Lee SK, Lui K, Yang J, Shah P; International Network for Evaluating Outcomes of Neonates (iNeo) Investigators. Neonatal outcomes of extremely preterm twins by sex pairing: an international cohort study. *Arch Dis Child Fetal Neonatal Ed.* 2020:fetalneonatal-2020-318832.
 64. Ushida T, Kotani T, Sadachi R, Hirakawa A, Hayakawa M, Moriyama Y, Imai K, Nakano-Kobayashi T, Kikkawa F; Neonatal Research Network of Japan. Antenatal Corticosteroids and Outcomes in Preterm Twins. *Obstet Gynecol.* 2020;135:1387-1397
 65. Ushida T, Kotani T, Hayakawa M, Hirakawa A, Sadachi R, Nakamura N, Moriyama Y, Imai K, Nakano-Kobayashi T, Kikkawa F. Antenatal corticosteroids and preterm offspring outcomes in hypertensive disorders of pregnancy: A Japanese cohort study. *Sci Rep.* 2020;10:9312.
 66. Arai H, Ito M, Ito T, Ota S, Takahashi T; Neonatal Research Network of Japan. Bubbly and cystic appearance on chest radiograph of extremely preterm infants with bronchopulmonary dysplasia is associated with wheezing disorder. *Acta Paediatr.* 2020;109:711-719.
 67. Hirata K, Kimura T, Hirano S, Wada K, Kusuda S, Fujimura M; Neonatal Research Network of Japan. Outcomes of outborn very-low-birth-weight infants in Japan. *Arch Dis Child Fetal Neonatal Ed.* 2020 Aug 11:fetalneonatal-2019-318594.
 68. Lehtonen L, Lee SK, Kusuda S, Lui K, Norman M, Bassler D, Håkansson S, Vento M, Darlow BA, Adams M, Puglia M, Isayama T, Noguchi A, Morisaki N, Helenius K, Reichman B, Shah PS; International Network for Evaluating Outcomes of Neonates (iNeo). Family Rooms in Neonatal Intensive Care Units and Neonatal Outcomes: An International Survey and Linked Cohort Study. *J Pediatr.* 2020:S0022-3476(20)30710-1
 69. Kurata H, Ochiai M, Inoue H, Ichiyama M, Yasuoka K, Fujiyoshi J, Matsushita Y, Honjo S, Sakai Y, Ohga S; Neonatal Research Network of Japan. A nationwide survey on tracheostomy for very-low-birth-weight infants in Japan. *Pediatr Pulmonol.* 2019;54:53-60.
 70. Lui K, Lee SK, Kusuda S, Adams M, Vento M, Reichman B, Darlow BA, Lehtonen L, Modi N, Norman M, Håkansson S, Bassler D, Rusconi F, Lodha A, Yang J, Shah PS; International Network for Evaluation of

- Outcomes (iNeo) of neonates Investigators. Trends in Outcomes for Neonates Born Very Preterm and Very Low Birth Weight in 11 High-Income Countries. *J Pediatr*.2019;215:32-40.e14.
71. Shahroor M, Lehtonen L, Lee SK, Håkansson S, Vento M, Darlow BA, Adams M, Mori A, Lui K, Bassler D, Morisaki N, Modi N, Noguchi A, Kusuda S, Beltempo M, Helenius K, Isayama T, Reichman B, Shah PS; on behalf of the International Network for Evaluation of Outcomes (iNeo) of neonates. Unit-Level Variations in Healthcare Professionals' Availability for Preterm Neonates <29 Weeks' Gestation: An International Survey. *Neonatology*. 2019;116:347-355.
 72. Matsushita Y, Sakai Y, Torio M, Inoue H, Ochiai M, Yasuoka K, Kurata H, Fujiyoshi J, Ichiyama M, Taguchi T, Kato K, Ohga S; Neonatal Research Network of Japan (NRNJ). Association of perinatal factors of epilepsy in very low birth weight infants, using a nationwide database in Japan. *J Perinatol*. 2019;39:1472-1479.
 73. Hitaka D, Morisaki N, Miyazono Y, Piedvache A, Nagafuji M, Takeuchi S, Kajikawa D, Kanai Y, Saito M, Takada H. Neonatal outcomes of very low birthweight infants born to mothers with hyperglycaemia in pregnancy: a retrospective cohort study in Japan. *BMJ Paediatr Open*. 2019;3:e000491.
 74. Yasuoka K, Inoue H, Egami N, Ochiai M, Tanaka K, Sawano T, Kurata H, Ichiyama M, Fujiyoshi J, Matsushita Y, Sakai Y, Ohga S; Neonatal Research Network of Japan. Late-Onset Circulatory Collapse and Risk of Cerebral Palsy in Extremely Preterm Infants. *J Pediatr*. 2019;212:117-123.e4.
 75. Morita M, Tanaka K, Matsumura S, Tamura M, Namba F. Perinatal factors associated with bubbly/cystic appearance in bronchopulmonary dysplasia: a nationwide, population-based cohort study in Japan. *J Matern Fetal Neonatal Med*. 2019;1-6.
 76. Arai H, Ito T, Ito M, Ota S, Takahashi T; Neonatal Research Network of Japan. Impact of chest radiography-based definition of bronchopulmonary dysplasia. *Pediatr Int*. 2019;61:258-263.
 77. Isayama T. The clinical management and outcomes of extremely preterm infants in Japan: past, present, and future. *Transl Pediatr*. 2019;8:199-211.
 78. Kusuda S. More studies are needed to convince stakeholders of the value of resuscitating periviable infants. *Acta Paediatr*. 2019;108:1182-1183.
 79. Inoue H, Ochiai M, Sakai Y, Yasuoka K, Tanaka K, Ichiyama M, Kurata H, Fujiyoshi J, Matsushita Y, Honjo S, Nonaka K, Taguchi T, Kato K, Ohga S; Neonatal Research Network of Japan. Neurodevelopmental Outcomes in Infants With Birth Weight ≤ 500 g at 3 Years of Age. *Pediatrics*. 2018;142:e20174286.
 80. Ichinomiya K, Maruyama K, Koizumi A, Inoue F, Fukuda K, Kaburagi K, Miyakawa Y; Neonatal Research Network of Japan. Comparison of

- neurodevelopmental outcomes between monochorionic and dichorionic twins with birth weight ≤ 1500 g in Japan: a register-based cohort study. *J Perinatol.* 2018 Oct;38(10):1407-1413.
81. Shah PS, Kusuda S, Håkansson S, Reichman B, Lui K, Lehtonen L, Modi N, Vento M, Adams M, Rusconi F, Norman M, Darlow BA, Lodha A, Yang J, Bassler D, Helenius KK, Isayama T, Lee SK; International Network for Evaluation of Outcomes (iNEO). Neonatal Outcomes of Very Preterm or Very Low Birth Weight Triplets. *Pediatrics.* 2018:e20181938
 82. Costeloe K, Turner MA, Padula MA, Shah PS, Modi N, Soll R, Haumont D, Kusuda S, Göpel W, Chang YS, Smith PB, Lui K, Davis JM, Hudson LD; International Neonatal Consortium. Sharing Data to Accelerate Medicine Development and Improve Neonatal Care: Data Standards and Harmonized Definitions. *J Pediatr* 2018; 203:437-441.
 83. Suzuki Y, Kono Y, Hayakawa T, Shimozawa H, Matano M, Yada Y; Neonatal Research Network Japan. Neonatal factors related to center variation in the incidence of late-onset circulatory collapse in extremely preterm infants. *PLoS One.* 2018;13:e0198518
 84. Darlow BA, Vento M, Beltempo M, Lehtonen L, Håkansson S, Reichman B, Helenius K, Sjörs G, Sigali E, Lee S, Noguchi A, Morisaki N, Kusuda S, Bassler D, San Feliciano L, Adams M, Isayama T, Shah PS, Lui K; on behalf of the International Network for Evaluating Outcomes (iNeo) of Neonates. Variations in Oxygen Saturation Targeting, and Retinopathy of Prematurity Screening and Treatment Criteria in Neonatal Intensive Care Units: An International Survey. *Neonatology* 2018;114:323-331.
 85. Persson M, Shah PS, Rusconi F, Reichman B, Modi N, Kusuda S, Lehtonen L, Håkansson S, Yang J, Isayama T, Beltempo M, Lee S, Norman M; International Network for Evaluating Outcomes of Neonates. Association of Maternal Diabetes With Neonatal Outcomes of Very Preterm and Very Low-Birth-Weight Infants: An International Cohort Study. *JAMA Pediatr.* 2018;172:867-875
 86. Totsu S, Terahara M, Kusuda S. Probiotics and the development of very low birthweight infants: follow-up study of a randomised trial. *BMJ Paediatr Open.* 2018;2(1):e000256
 87. Nakanishi H, Suenaga H, Uchiyama A, Kono Y, Kusuda S; Neonatal Research Network, Japan. Trends in the neurodevelopmental outcomes among preterm infants from 2003-2012: a retrospective cohort study in Japan. *J Perinatol.* 2018;38:917-928.
 88. Beltempo M, Isayama T, Vento M, Lui K, Kusuda S, Lehtonen L, Sjörs G, Håkansson S, Adams M, Noguchi A, Reichman B, Darlow BA, Morisaki N, Bassler D, Pratesi S, Lee SK, Lodha A, Modi N, Helenius K, Shah PS; on behalf of the International Network for Evaluating Outcomes of Neonates. Respiratory Management of Extremely Preterm Infants: An International Survey. *Neonatology.* 2018;114(1):28-36
 89. Kono Y, Yonemoto N, Nakanishi H, Kusuda S, Fujimura M. Changes in

- survival and neurodevelopmental outcomes of infants born at <25 weeks' gestation: a retrospective observational study in tertiary centres in Japan. *BMJ Paediatr Open*. 2018;2(1):e000211
90. Nakanishi H, Suenaga H, Uchiyama A, Kusuda S; Neonatal Research Network, Japan. Persistent pulmonary hypertension of the newborn in extremely preterm infants: a Japanese cohort study. *Arch Dis Child Fetal Neonatal Ed*. 2018;103:F554-561.
 91. Morisaki N, Isayama T, Samura O, Wada K, Kusuda S. Socioeconomic inequity in survival for deliveries at 22-24 weeks of gestation. *Arch Dis Child Fetal Neonatal Ed*. 2018;103:F202-F207
 92. Helenius K, Sjörs G, Shah PS, Modi N, Reichman B, Morisaki N, Kusuda S, Lui K, Darlow BA, Bassler D, Håkansson S, Adams M, Vento M, Rusconi F, Isayama T, Lee SK, Lehtonen L; International Network for Evaluating Outcomes (iNeo) of Neonates. Survival in Very Preterm Infants: An International Comparison of 10 National Neonatal Networks. *Pediatrics*. 2017 2017 ;140: e20171264
 93. Inoue H, Ochiai M, Yasuoka K, Tanaka K, Kurata H, Fujiyoshi J, Matsushita Y, Suga S, Nonaka K, Taguchi T, Kato K, Ohga S; Neonatal Research Network of Japan (NRNJ). Early Mortality and Morbidity in Infants with Birth Weight of 500 Grams or Less in Japan. *J Pediatr*. 2017;190:112-117.e3.
 94. Vento M, Rusconi F, Lehtonen L, Reichman B, Darlow BA, Lui K, Feliciano LS, Gagliardi L, Bassler D, Modi N, Kelly LE, Shah PS, Håkansson S, Kusuda S, Adams M, Lee SK, Sjörs G, Perinatal health services organization for preterm births: a multinational comparison. *J Perinatol*. 2017;37:762-768
 95. Kimura T, Takeuchi M, Imai T, Tanaka S, Kawakami K; Neonatal Research Network of Japan. Neurodevelopment at 3 Years in Neonates Born by Vaginal Delivery versus Cesarean Section at <26 Weeks of Gestation: Retrospective Analysis of a Nationwide Registry in Japan. *Neonatology*. 2017;112:258-266.
 96. Ito M, Tamura M, Namba F; Neonatal Research Network of Japan. Role of sex in morbidity and mortality of very premature neonates. *Pediatr Int*. 2017;59:898-905.
 97. Sasaki H, Yonemoto N, Mori R, Nishida T, Kusuda S, Nakayama T. Assessing archetypes of organizational culture based on the Competing Values Framework: the experimental use of the framework in Japanese neonatal intensive care units. *Int J Qual Health Care*. 2017;29:384-391.
 98. Darlow BA, Lui K, Kusuda S, Reichman B, Håkansson S, Bassler D, Modi N, Lee SK, Lehtonen L, Vento M, Isayama T, Sjörs G, Helenius KK, Adams M, Rusconi F, Morisaki N, Shah PS; International Network for Evaluating Outcomes of Neonates. International variations and trends in the treatment for retinopathy of prematurity. *Br J Ophthalmol*. 2017;101:1399-1404
 99. Hines D, Modi N, Lee SK, Isayama T, Sjörs G, Gagliardi L, Lehtonen L,

- Vento M, Kusuda S, Bassler D, Mori R, Reichman B, Håkansson S, Darlow BA, Adams M, Rusconi F, San Feliciano L, Lui K, Morisaki N, Musrap N, Shah PS; International Network for Evaluating Outcomes (iNeo) of Neonates.. Scoping review shows wide variation in the definitions of bronchopulmonary dysplasia in preterm infants and calls for a consensus. *Acta Paediatr.* 2017;106:366-374.
100. Gemmell L, Martin L, Murphy KE, Modi N, Håkansson S, Reichman B, Lui K, Kusuda S, Sjörs G, Mirea L, Darlow BA, Mori R, Lee SK, Shah PS. Hypertensive disorders of pregnancy and outcomes of preterm infants of 24 to 28 weeks' gestation. *J Perinatol.* 2016;36:1067-1072.
 101. Nakanishi H, Uchiyama A, Kusuda S. Impact of pulmonary hypertension on neurodevelopmental outcome in preterm infants with bronchopulmonary dysplasia: a cohort study. *J Perinatol.* 2016;36:890-6.
 102. Shah PS, Lui K, Sjörs G, Mirea L, Reichman B, Adams M, Modi N, Darlow BA, Kusuda S, San Feliciano L, Yang J, Håkansson S, Mori R, Bassler D, Figueras-Aloy J, Lee SK. Neonatal Outcomes of Very Low Birth Weight and Very Preterm Neonates: An International Comparison.; International Network for Evaluating Outcomes (iNeo) of Neonates. *J Pediatr.* 2016;177:144-152.e6
 103. Martin LJ, Sjörs G, Reichman B, Darlow BA, Morisaki N, Modi N, Bassler D, Mirea L, Adams M, Kusuda S, Lui K, Feliciano LS, Håkansson S, Isayama T, Mori R, Vento M, Lee SK, Shah PS; International Network for Evaluating Outcomes (iNeo) of Neonates Investigators. Country-Specific vs. Common Birthweight-for-Gestational Age References to Identify Small for Gestational Age Infants Born at 24-28 weeks: An International Study. *Paediatr Perinat Epidemiol.* 2016;30:450-61.
 104. Nakamura T, Yonemoto N, Nakayama M, Hirano S, Aotani H, Kusuda S, Fujimura M, Tamura M; and The Neonatal Research Network, Japan. Early inhaled steroid use in extremely low birthweight infants: a randomised controlled trial. *Arch Dis Child Fetal Neonatal Ed.* 2016;10:F552-F556.
 105. Ohhashi M, Yoshitomi T, Sumiyoshi K, Kawagoe Y, Satoh S, Sameshima H, Ikenoue T. Magnesium Sulfate and perinatal mortality and morbidity in very-low-birthweight infants born between 24 and 32 weeks of gestation in Japan. *Eur J Obstet Gynecol Reprod Biol.* 2016;201:140-145.
 106. Yamakawa T, Itabashi K, Kusuda S; Neonatal Research Network of Japan. Mortality and morbidity risks vary with birth weight standard deviation score in growth restricted extremely preterm infants. *Early Hum Dev* 2016;92:7-11.
 107. Miyazaki K, Furuhashi M, Ishikawa K, Tamakoshi K, Hayashi K, Kai A, Ishikawa H, Murabayashi N, Ikeda T, Kono Y, Kusuda S, Fujimura M. Impact of chorioamnionitis on short- and long-term outcomes in very low birth weight preterm infants: the Neonatal

- Research Network Japan. *J Matern Fetal Neonatal Med.* 2016;29:331-7.
108. Miyazaki K, Furuhashi M, Ishikawa K, Tamakoshi K, Hayashi K, Kai A, Ishikawa H, Murabayashi N, Ikeda T, Kono Y, Kusuda S, Fujimura M. Long-term outcomes of antenatal corticosteroids treatment in very preterm infants after chorioamnionitis. *Arch Gynecol Obstet.* 2015;292:1239-46.
 109. Isayama T, Mirea L, Mori R, Kusuda S, Fujimura M, Lee SK, Shah PS; Neonatal Research Network of Japan and the Canadian Neonatal Network. Patent ductus arteriosus management and outcomes in Japan and Canada: comparison of proactive and selective approaches. *Am J Perinatol.* 2015;32:1087-94.
 110. Isayama T, Ye XY, Tokumasu H, Chiba H, Mitsuhashi H, Shahrook S, Kusuda S, Fujimura M, Toyoshima K, Mori R; Neonatal Research Network of Japan. The effect of professional-led guideline workshops on clinical practice for the management of patent ductus arteriosus in preterm neonates in Japan: a controlled before-and-after study. *Implement Sci.* 2015;10:67.
 111. Ishikawa H, Miyazaki K, Ikeda T, Murabayashi N, Hayashi K, Kai A, Ishikawa K, Miyamoto Y, Nishimura K, Kono Y, Kusuda S, Fujimura M; Neonatal Research Network of Japan. The Effects of Antenatal Corticosteroids on Short- and Long-Term Outcomes in Small-for-Gestational-Age Infants. *Int J Med Sci.* 2015;12(4):295-300.
 112. Maruyama H, Yonemoto N, Kono Y, Kusuda S, Fujimura M; Neonatal Research Network of Japan. Weight Growth Velocity and Neurodevelopmental Outcomes in Extremely Low Birth Weight Infants. *PLoS One.* 2015;10:e0139014.
 113. Totsu S, Yamasaki C, Terahara M, Uchiyama A, Kusuda S; Probiotics Study Group in Japan. Bifidobacterium and enteral feeding in preterm infants: cluster-randomized trial. *Pediatr Int* 2014;56:714-9.
 114. Miyazaki K, Furuhashi M, Ishikawa K, Tamakoshi K, Ikeda T, Kusuda S, Fujimura M; Neonatal Research Network Japan. The effects of antenatal corticosteroids therapy on very preterm infants after chorioamnionitis. *Arch Gynecol Obstet.* 2014;289:1185-90.
 115. Shah PS, Lee SK, Lui K, Sjörs G, Mori R, Reichman B, Håkansson S, Feliciano LS, Modi N, Adams M, Darlow B, Fujimura M, Kusuda S, Haslam R, Mirea L; International Network for Evaluating Outcomes of Neonates (iNeo). The International Network for Evaluating Outcomes of very low birth weight, very preterm neonates (iNeo): a protocol for collaborative comparisons of international health services for quality improvement in neonatal care. *BMC Pediatr.* 2014;14:110.
 116. Sasaki Y, Ikeda T, Nishimura K, Katsuragi S, Sengoku K, Kusuda S, Fujimura M. Association of antenatal corticosteroids and the mode of delivery with the mortality and morbidity of infants weighing less than 1,500g at birth in Japan. *Neonatology.* 2014;106:81-6.
 117. Morisaki N, Belfort MB, McCormick MC, Mori R, Noma H, Kusuda

- S, Fujimura M; Neonatal Research Network of Japan. Brief parenteral nutrition accelerates weight gain, head growth even in healthy VLBWs. *PLoS One* 2014;9:e88392.
118. Kusuda S, Fujimura M, Uchiyama A, Nakanishi H, Totsu S; for Neonatal Research Network, Japan. Identification of practices and morbidities affecting the mortality of very low birth weight infants using a multilevel logistic analysis: clinical trial or standardization? *BMJ Open* 2013;3:e003317.
 119. Ishii N, Kono Y, Yonemoto N, Kusuda S, Fujimura M; Neonatal Research Network, Japan. Outcomes of infants born at 22 and 23 weeks' gestation. *Pediatrics*. 2013;132:62-71.
 120. Kusuda S, Fujimura M, Uchiyama A, Totsu S, Matsunami K; Neonatal Research Network, Japan. Trends in morbidity and mortality among very-low-birth-weight infants from 2003 to 2008 in Japan. *Pediatr Res* 2012;72:531-8.
 121. Isayama T, Lee SK, Mori R, Kusuda S, Fujimura M, Ye XY, Shah PS; Canadian Neonatal Network; Neonatal Research Network of Japan. Comparison of mortality and morbidity of very low birth weight infants between Canada and Japan. *Pediatrics* 2012;130:e957-65.
 122. Kono Y, Mishina J, Yonemoto N, Kusuda S, Fujimura M. Outcomes of very-low-birthweight infants at 3 years of age born in 2003-2004 in Japan. *Pediatr Int* 2011;53:1051-8.
 123. Kono Y, Mishina J, Yonemoto N, Kusuda S, Fujimura M; NICU Network, Japan. Neonatal correlates of adverse outcomes in very low-birthweight infants in the NICU Network. *Pediatr Int* 2011;53:930-5.
 124. Mori R, Kusuda S, Fujimura M; Neonatal Research Network Japan. Antenatal corticosteroids promote survival of extremely preterm infants born at 22 to 23 weeks of gestation. *J Pediatr* 2011;159:110-114.
 125. Valls-I-Soler A, Azpeitia A, Madrid M, Kusuda S, Mori M; Spanish Neonatal Network (SEN1500) & Neonatal Research Network of Japan (NRNJ). Neonatal Mortality Rate (NMR) in Very Low Birth Weight Infants (VLBWI). Comparison Between Neonatal Networks from Spain (SEN1500) and Japan (NRNJ). *Pediatr Res* 2011;70:36.
 126. Kusuda S, Fujimura M, Sakuma I, Aotani H, Kabe K, Itani Y, Ichiba H, Matsunami K, Nishida H; Neonatal Research Network, Japan. Morbidity and mortality of infants with very low birth weight in Japan: center variation. *Pediatrics* 2006;118:e1130-8.